

## XVIII. SOME BATS FROM OKLAHOMA.\*

### A. I. Ortenburger.

The bats listed below, with a single exception, were collected during the summer of 1925. It is interesting that four different genera are represented by the relatively few specimens which it was possible to secure. The author is indebted to the Bureau of Biological Survey, U. S. Department of Agriculture, for checking the identifications.

#### *Pipistrellus subflavus subflavus* (F. Cuvier).

U. O. M. Z. 2103, 2104, 2105. McCurtain Co., 10 miles southeast of Broken Bow. Shot while flying over Mountain Fork River.

U. O. M. Z. 2107, 2257. McCurtain County, 1 mile west of the Oklahoma-Arkansas line on the Red River. Shot while flying over the Red River. This form as well as others not obtained was very numerous.

#### *Nyctinomus mexicanus* (Bailey).

U. O. M. Z. 2108. Cleveland Co., Norman, on the University of Oklahoma campus in the Law Building.

#### *Nycteris borealis borealis* (Muller).

U. O. M. Z. 2106. McCurtain County, 10 miles southeast of Broken Bow. Shot while flying over Mountain Fork River.

#### *Nycticeius humeralis* (Raf.).

U. O. M. Z. 2101, 2102. Pushmataha County, 4 miles northeast of Tuskahoma. Shot while flying over the Kiamichi River. Very common.

---

\*Contribution from the Zoological Laboratory of the University of Oklahoma, Second Series, No. 65.