

## A List of Mammals Collected in West Central Oklahoma

OTIS M. KING, Southwestern State College, Weatherford, Oklahoma

The mammals comprising the following list were collected since November, 1950. Most of the collections were made in Custer County; however, some locality records are included from Alfalfa, Blaine, Caddo, Kiowa, Major, and Washita Counties. Sight records are not included. Skins or skulls have been prepared for each specimen.

**ACKNOWLEDGEMENTS:** Specimens were identified at the University of Kansas Museum of Natural History by Dr. Rollin H. Baker, now of Michigan State University, and Mr. Robert J. Russell, now of the University of Kansas City. Dr. Bryan P. Glass of Oklahoma A. and M. College also identified some specimens.

### Annotated List

*Didelphis virginiana virginiana* Kerr. Blaine Co.: 10 miles north, 7 miles east of Weatherford. Custer Co.: 2 miles east of Weatherford; 1½ miles west of Weatherford.

*Scalopus aquaticus intermedius* (Elliot). Custer Co.: Weatherford.

*Cryptotis parva parva* (Say). Custer Co.: 1½ miles north, ¾ miles west of weatherford.

*Myotis velifer incautus* (J. A. Allen). Custer Co.: 3 miles north, 6 miles west of Weatherford; 3 miles east of Weatherford; Butler. Washita Co.: 5 miles south, 5 miles west of Weatherford; 5 miles west, 1 mile south of Foss.

*Pipistrellus subflavus subflavus* (F. Cuvier). Washita Co.: 5 miles south, 5 miles west of Weatherford.

*Lasturus borealis borealis* (Muller). Custer Co.: Weatherford; 4 miles northwest of Weatherford.

*Corynorhinus rafinesquii pallescens* Miller. Washita Co.: 5 miles south, 5 miles west of Weatherford; 4 miles west, 6 miles south of Weatherford.

*Tadarida brasiliensis mexicana* (Saussure). Custer Co.: Weatherford.

*Procyon lotor hirtus* Nelson and Goldman. Custer Co.: 2½ miles southwest of Thomas; 8 miles north, 4 miles east of Weatherford; 5 miles north of Weatherford; 2 miles east of Weatherford.

*Spilogale putorius interrupta* (Rafinesque). Washita Co.: ¾ miles north, 3 miles east of Lake Valley. Kiowa Co.: ¾ miles south of Hobart.

*Mephitis mephitis* Gray. Custer Co.: Weatherford; 1½ miles east of Weatherford; 2 miles west of Butler.

*Taxidea taxus berlandieri* Baird. Custer Co.: 6 miles south of Thomas; 1½ miles east of Weatherford.

*Urocyon cinereoargenteus ocythous* Bangs. Custer Co.: ¾ miles south of Weatherford; 4 miles south of Weatherford. Caddo Co.: 8 miles west of Lookeba. Kiowa Co.: 3 miles north of Roosevelt.

*Canis latrans nebracensis* Merriam. Custer Co.: 1 mile west of Weatherford. Washita Co.: 9 miles west, 8 miles south of Weatherford. Kiowa Co.: 3 miles north of Hobart.

*Lynx rufus baileyi* (Merriam). Major Co.: 12 miles northeast of Selling.

*Citellus tridecemlineatus texensis* (Merriam). Custer Co.: 3 miles north, 3 miles east of Weatherford; 3 miles south of Weatherford.

*Oynomys ludovicianus ludovicianus* (Ord). Custer Co.: 15 miles north of Weatherford; 1 mile north, 2 miles west of Weatherford. Washita Co.: 7 miles south, 2 miles east of Weatherford.

*Sciurus niger rufiventer* Geoffrey. Blaine Co.: 10 miles north, 7 miles east of Weatherford. Custer Co.: 15 miles north of Weatherford; 5 miles north of Weatherford.

*Geomys bursarius major* Davis. Blaine Co.: 1½ miles north of Hydro. Custer Co.: Weatherford.

*Perognathus flavus* Baird. Custer Co.: 4 miles west, 4½ miles north of Weatherford.

*Perognathus hispidus paradoxus* Merriam. Blaine Co.: 1 mile north of Hydro. Custer Co.: 5 miles north, 4 miles west of Weatherford; 4 miles south, 2 miles west of Weatherford; 6 miles west of Weatherford.

*Dipodomys ordii richardsoni* (Allen). Blaine Co.: 2 miles north, 1 mile east of Hydro. Custer Co.: 4 miles south of Weatherford; 3 miles west of Butler. Washita Co.: 7 miles south, 2½ miles east of Weatherford.

*Onychomys leucogaster breviauritus* Hollister. Blaine Co.: 2 miles north, 1 mile west of Hydro. Custer Co.: 7 miles north, 5 miles west of Weatherford; ¾ mile north, 2 miles east of Weatherford; 4 miles south, 2 miles west of Weatherford; 11 miles west of Weatherford; 3 miles west of Butler.

*Reithrodontomys montanus griseus* Bailey. Custer Co.: 4 miles north, 2 miles west of Weatherford; 1 mile north, 2½ miles west of Weatherford; 1 mile south, 5 miles west of Weatherford.

*Peromyscus maniculatus nebrascensis* (Coues). Blaine Co.: 2 miles north of Hydro. Custer Co.: 7 miles north, 5 miles west of Weatherford; 4 miles north, 2 miles west of Weatherford; 11 miles west of Weatherford; 7 miles west of Butler. Washita Co.: 4½ miles south of Weatherford.

*Peromyscus leucopus leucopus* (Rafinesque). Blaine Co.: 17 miles south-west of Okeene. Custer Co.: 5 miles north of Weatherford; 1½ miles east, ½ mile south of Weatherford; 7 miles west of Weatherford; 7 miles west of Butler. Washita Co.: 4½ miles south of Weatherford.

*Sigmodon hispidus texianus* (Audubon and Bachman). Blaine Co.: 2 miles north, 1 mile west of Hydro. Custer Co.: 5 miles north of Weatherford; 1½ miles south, 1 mile west of Weatherford; 4 miles east of Butler.

*Neotoma micropus micropus* Baird. Custer Co.: 2 miles north, 1.3 miles west of Weatherford.

*Ondatra zibethicus cinnamominus* (Hollister). Alfalfa Co.: 15 miles north of Jet.

*Rattus norvegicus* (Berkenhout). Custer Co.: Weatherford.

*Mus musculus* Linnaeus. Custer Co.: Weatherford.

*Lepus californicus melanotis* Mearns. Custer Co.: 1 mile north, 2 miles west of Weatherford; 1½ mile south, 2 miles west of Weatherford.

*Sylvilagus floridanus llanensis* Blair. Custer Co.: 15 miles north of Weatherford. Washita Co.: 7 miles south, 3 miles east of Weatherford.