
A New Geographic Record for *Brachycybe lecontii* (Diplopoda: Platydesmida: Andrognathidae) from Oklahoma

Chris T. McAllister

Science and Mathematics Division, Eastern Oklahoma State College, Idabel, OK 74745

Matthew B. Connior

Life Sciences, Northwest Arkansas Community College, One College Drive, Bentonville, AR 72712

©2016 Oklahoma Academy of Science

Much has been published on the millipeds (Diplopoda) of Oklahoma since the turn of the century (Lewis 2002; McAllister et al. 2002, 2005, 2013; McAllister and Shelley 2003a, 2003b, 2005, 2008, 2010; Shear 2003; Shelley et al. 2005a, 2005b, 2006; McAllister and Robison 2011; Lewis and Slay 2012; Shelley and Snyder 2012). However, there are still gaps in the distribution of many species. Here, we continue to help fill some of those gaps with a noteworthy county record of an uncommon milliped of the state. Additionally, we provide an updated distribution for this species in Oklahoma and surrounding states.

On 1 April 2016, MBC collected two millipeds from a cave/spring outflow site off county road 660 at Flint, Delaware County (36° 12' 27.83"N, 94° 42' 15.78"W). They were taken off the ground near the cave entrance. Specimens were placed in a vial containing 70% (v/v) ethanol and, following preliminary identification, were sent to CTM for verification of identification and deposition of voucher specimens in the Sam Noble Oklahoma Museum of Natural History (SNOMNH, Acc. No. 1579), Norman.

Based on morphological characters, the millipeds fit the description of *Brachycybe lecontii* Wood, 1864 using the key to species of *Brachycybe* of Gardner (1975), modified by Shelley et al. (2005b). Our specimens displayed a bright pinkish-red paranota which is typical

of *B. lecontii* in the western part of its range (McAllister et al. 2002). We did not record sexes because this milliped is the only representative of the Platydesmida in the state that can be authentically identified with either sex and/or with juveniles.

This milliped had been previously reported from only two counties, Le Flore and McCurtain (Appendix) in the southeastern corner of the state on the Ouachita uplift. It is typically found under decaying hardwood logs or under the bark of those logs but as yet not under pine logs or bark. The current location (Flint) is important as it is on the Ozark Plateau of Oklahoma, which extends northward and ends in southeastern Kansas in the vicinity of Galena, Cherokee County (see Shelley et al. 2005b, Fig. 19 showing the outlined projected range for "population 2" of *B. lecontii*). The potential Kansas location (Fig. 1) would be geographically important because the authors and others, have attempted to find *B. lecontii* there at Schermerhorn Park (3.7 km S of Galena off St. Hwy 26) in typical habitat (moist upland deciduous woods) on several occasions but were unsuccessful. The reason for these failures remain an enigma but we suggest continued surveying in this region as its collection would be a new state record for Kansas.

Interestingly, a recent molecular study of *Brachycybe* (Brewer et al. 2012) did not

recognize multiple species within the *B. lecontii* clade. In addition, “populations 2, 3” of Shelley et al. (2005b), which included specimens from Le Flore and McCurtain counties (Appendix), represented a genetically divergent lineage that shows considerable internal geographic structuring (Brewer et al. 2005).

Brachybybe lecontii has a large range that occupies a north to south area extending from southern West Virginia, central Kentucky, and southern Missouri to the southern periphery of Georgia, southcentral Louisiana, southeastern Texas, and possibly southeastern Kansas (Gardner 1975; McAllister et al. 2003b; Shelley et al. 2005b, see Figs. 18–19; Youngsteadt and McAllister 2014). In Arkansas, it occurs along the northwestern tier of counties (Benton, Crawford and Washington) along the Oklahoma state line (McAllister et al. 2003b) (Fig. 1). Here, we officially extend it into northeastern Oklahoma with deposition of genuine voucher specimens, which constitutes a new county record, and, most importantly, a noteworthy extension of the northwest range distribution of the species in North America.




Figure 1. Distribution of *Brachybybe lecontii* in Arkansas, Louisiana, Missouri, Oklahoma and Texas. Dots = previous records; star = new record. Arrow denotes potential occurrence in Cherokee County of southeastern Kansas.

Acknowledgments

The Oklahoma Department of Wildlife Conservation provided a scientific collecting permit to CTM. We thank Dr. Jason E. Bond (Auburn University) for assistance in locating some records.

References

- Black JH. 1971. The cave life of Oklahoma: A preliminary study (excluding Chiroptera). *Okla. Undergr.* 4:2–56.
- Brewer MS, Spruill CL, Rao NS, Bond JE. 2012. Phylogenetics of the millipede genus *Brachybybe* Wood, 1864 (Diplopoda: Platydesmida: Andrognathidae): patterns of deep evolutionary history and recent speciation. *Mol. Phylo. Evol.* 64:232–242.
- Gardner MR. 1975. Revision of the millipede family Andrognathidae in the Nearctic region (Diplopoda, Platydesmida). *Mem. Pac. Coast Entomol. Soc.* 5:1–61.
- Lewis JJ. 2002. *Chaetaspis aleyorum*, a new species of millipede from Tumbling Creek Cave, Missouri, with a synopsis of the cavernicolous species of *Chaetaspis* (Diplopoda: Polydesmida): *Myriapodologica* 7:101–111.
- Lewis JJ, Slay ME. 2012. *Chaetaspis attenuatus*, a new species of cavernicolous millipede from Arkansas (Diplopoda: Polydesmida: Macrosterodesmidae). *J. Cave Karst Stud.* 75:60–63.
- McAllister CT, Robison HW. 2011. Millipeds (Arthropoda: Diplopoda) of the Ark-La-Tex. V. New distribution records for select taxa in Arkansas, Oklahoma, and Texas. *Southwest. Nat.* 56:422–426.
- McAllister CT, Robison HW, Connior MB. 2013. New geographic distribution records for millipeds (Diplopoda) from Oklahoma. *Proc. Okla. Acad. Sci.* 93:7–12.
- McAllister CT, Shelley RM. 2003a. Millipeds (Arthropoda: Diplopoda) of the Ark-La-Tex. IV. New geographic distribution records from southcentral and southeastern Oklahoma. *J. Ark. Acad. Sci.* 57:115–121.

- McAllister CT, Shelley RM. 2005. Discovery of the milliped, *Tiganogona brownae* Chamberlin (Diplopoda: Chordeumatida: Cleidogonidae) in central Oklahoma: westernmost records for the family and genus. *Proc. Okla. Acad. Sci.* 85:79–80.
- McAllister CT, Shelley RM. 2008. New records of eurymerodesmid millipeds (Diplopoda: Polydesmida) from Arkansas, Kansas, Louisiana, Oklahoma, and Texas. *J. Ark. Acad. Sci.* 62:155–158.
- McAllister CT, Shelley RM. 2010. Distribution of *Abacion texense* (Loomis, 1937), the only milliped species traversing the Rio Grande, Mississippi, and Pecos rivers (Callipodida: Abacionidae). *Insecta Mundi* 124:1–8.
- McAllister CT, Shelley RM, Enghoff H, Ramsey ZD. 2005. Distribution of the milliped *Virgoiulus minutus* (Brandt, 1841): first records from Mississippi, Oklahoma, and Texas (Julida: Blaniulidae). *West. North Amer. Nat.* 65:258–266.
- McAllister CT, Shelley RM, McAllister JT III. 2002. Millipeds (Arthropoda: Diplopoda) of the Ark-La-Tex. II. Distributional records for some species of western and central Arkansas and eastern and southeastern Oklahoma. *J. Ark. Acad. Sci.* 56:95–98.
- McAllister CT, Shelley RM, McAllister JT III. 2003b. Millipeds (Arthropoda: Diplopoda) of the Ark-La-Tex. III. Additional records from Arkansas. *J. Ark. Acad. Sci.* 57:95–98.
- Shear WA. 2003. The milliped family Trichopetalidae, Part 1: Introduction and genera *Trigenotyla* Causey, *Nannopetalum* n. gen., and *Causeyella* n. gen. (Diplopoda: Chordeumatida, Cleidogonoidea). *Zootaxa* 321:1–36.
- Shelley RM, McAllister CT, Medrano MF. 2006. Distribution of the milliped genus *Narceus* Rafinesque, 1820 (Spirobolida: Spirobolidae): occurrences in New England and west of the Mississippi River: a summary of peripheral localities; and first records from Connecticut, Delaware, Maine, and Minnesota. *West. North Amer. Nat.* 66:374–389.
- Shelley RM, McAllister, Ramsey ZD. 2005a. Discovery of the milliped *Scytonotus granulatus* (Say, 1821) in Oklahoma and Alabama, with a review of its distribution (Polydesmida: Polydesmidae). *West. North Amer. Nat.* 65:112–117.
- Shelley RM, McAllister CT, Tanabe T. 2005b. A synopsis of the milliped genus *Brachycybe* Wood, 1864 (Platydesmida: Andrognathidae). *Frag. Faun.* 48:137–166.
- Shelley RM, Snyder BA. 2012. Millipeds of the eastern Dakotas and western Minnesota, USA, with an account of *Pseudopolydesmus serratus* (Say, 1821) (Polydesmida: Polydesmidae); first published records from six states and the District of Columbia. *Insecta Mundi* 239:1–17.
- Youngsteadt NW, McAllister CT. 2014. Natural history notes and new county records for Ozarkian millipeds (Arthropoda: Diplopoda) from Arkansas, Kansas and Missouri. *J. Ark. Acad. Sci.* 68:177–182.

Submitted August 9, 2016 Accepted November 21, 2016

Appendix. Summary of collection data on *Brachycybe lecontii* from Oklahoma.

Delaware County

(1) Vicinity of Flint off county road 660, 1 Apr. 2016, M.B. Connior (SNOMNH).

Le Flore County

(1) 11.2 km NE Talihina at Bear Den Cave, 29 Apr. 1971, J.H. Black (FSCA) (see Black 1971).

(2) Choctaw Nation Historic site at junction of St. Hwy 1 and AR St. Hwy 88, 27 Sept. 2002 & 13 Aug. 2003, C.T. McAllister (NCSM).

(3) Off US 259, ~9.7 km N of St. Hwy 144 junction. No date. J.E. Bond (AUM).

(4) Ouachita National Forest on US 59, ~12.9 km E of US 259 junction near Oklahoma welcome sign. No date. J.E. Bond (AUM).

McCurtain County

(1) Beavers Bend State Park, 16 Jun. 1970, 11 June 1985, D.C. Arnold (OKSU).

(2) Beavers Bend State Park, David Boren Trail, 29 Dec. 2003, C.T. & R.A. McAllister (NCSM).

(3) 4.8 km N of Tom, 7 Feb. 2004, Z.D. Ramsey (NCSM).

(4) Halibut Bay Road, Hochatown. 15 Nov. 2015, C.T. McAllister (author's collection).

(5) Off US 259, ~9.7 km SW of St. Hwy 4 junction. No date. J.E. Bond (AUM).

Museum acronyms: AUM = Auburn University Museum of Natural History, Auburn, Alabama; FSCA = Florida State Collection of Arthropods, Gainesville; NCSM = North Carolina State Museum of Natural Sciences, Raleigh; OKSU = Department of Entomology and Plant Pathology, Oklahoma State University, Stillwater; SNOMNH = Sam Noble Oklahoma Museum of Natural History, Norman.

Age and Size of Spotted Gar (*Lepisosteus oculatus*) from Lake Thunderbird Reservoir in Central Oklahoma

Bryan D. Frenette

Department of Biology, University of Oklahoma, Norman, OK 73019

Richard A. Snow

Oklahoma Department of Wildlife Conservation, Oklahoma Fishery Research Lab,
Norman, OK 73072

Abstract: We report the body size and population age structure of Spotted Gar from the Lake Thunderbird reservoir, OK. We collected 90 Spotted Gars ranging between 348–846 mm total length, and aged gars by examining annuli from the sagittal otoliths. Annuli were counted on both browned and not browned sagittal otoliths to compare the efficacy of both methods. Gars ranged between 1 and 14 years old, and percent agreement of age within one year between readers was comparable for both the browned and not browned otoliths. Back-calculated growth indicated that Spotted Gar grow quickly during the first year of life, and approach maximum size by year 4. ©2016 Oklahoma Academy of Science

Introduction

The Spotted Gar (*Lepisosteus oculatus*) is one of four species of gar (Family: Lepisosteidae) that occur in Oklahoma. While globally secure, the Spotted Gar is a species of conservation concern at the northern edge of its range and is critically imperiled in Canada (Glass et al. 2011; Statton et al. 2012; David et al. 2015; NatureServe 2016), critically imperiled in Kansas, Ohio, and Pennsylvania, and is thought to be extirpated in New Mexico (Staton et al. 2012; NatureServe 2016). The basic biology of this species and other gars remains largely understudied, due in part to the reputation gars hold as nuisance fish throughout much of their range (Scarnecchia 1992). This lack of information concerning Spotted Gar holds true for Oklahoma, where few studies have focused on this species. Echelle and Riggs (1972) described early life history of gars, including Spotted Gar, in Lake Texoma. Tyler and Granger (1984) reported on size, diet, and spawning behavior of Spotted Gar from Lake Lawtonka, and Frenette and Snow (2016)

described spawning behavior in Spotted Gar from Lake Thunderbird. This study describes population age using sagittal otoliths and body size of Spotted Gar in the Lake Thunderbird reservoir.

Methods

Study Area Description

The Lake Thunderbird reservoir is a man-made impoundment located in the Cleveland and Oklahoma counties in central Oklahoma, USA (Figure 1; Simonds 1999). The reservoir was built and put into operation between 1962 and 1965, and is located approximately 16km from the city of Norman, OK (Simonds 1999). Lake Thunderbird impounds the Little River, a tributary to the Canadian River, and provides water for Oklahoma City, Norman, Midwest City, Del City, and the Tinker Air Force Base (Simonds 1999). Historic and contemporary sampling did not detect Spotted Gars or other species of Lepisosteidae in the Little River and its tributaries surrounding Lake Thunderbird (Franssen and Tobler 2013). However, sampling methods from this study may not have been




Figure 1.—Map of the Lake Thunderbird reservoir in central Oklahoma.

sufficient to efficiently capture large-bodied fishes like Gars (Franssen and Tobler 2013; M. Tobler, Kansas State University, personal communication). It was hypothesized that flooding in the late 1990's allowed Spotted Gar to migrate into Lake Thunderbird from Lake Stanley Draper to the north; Spotted Gar were not detected by sampling in Lake Thunderbird until the mid-2000's (K. Thomas, Oklahoma Department of Wildlife Conservation, personal communication).

Sampling and Data Collection

Sampling on April 25 and May 2, 2014 in the Hog Creek arm of Lake Thunderbird captured 90 adult Spotted Gar. Collections were made using boat electroshocking in areas where gar densities were high, presumably due to the spawning season. Fish were transported to the Aquatic Research Facility (ARF) at the University of Oklahoma, where they were held in a 0.04 ha holding pond until being processed for this study.

Sex identification and morphometric measures.—Prior to measuring, fish were

placed into an ice bath to induce torpor. Fish were measured in mm for total length (TL) using a measuring board (± 1 mm). Calipers were used to measure both head length (HL) and snout length (SnL) in mm. HL was measured from the tip of the snout to the posterior portion of the opercle. SnL was measured from the tip of the snout to the anterior start of the orbital. Both HL and SnL were measured alongside the snout on the left side of the body. Fish were weighed in kg using a digital scale. Sex of Spotted Gars was determined by examining the gamete release pathways following a standardized procedure for determining sex in Lepisosteids (Ferrara and Irwin 2001).

Otolith aging.—Paired sagittal otoliths were removed from Spotted Gar through the ventral side of the brain case. Otoliths were then cleaned and dried before processing. One otolith from each pair was browned at 104°C using a hot plate to increase the contrast between the accretion and discontinuous zones (Figure 2; Secor et al. 1992; Long and Snow 2016). Both otoliths (browned and not browned) were processed following Buckmeier et al. (2012),




Figure 2.—Whole (WH) sagittal otolith view to the left with not browned (NB) and browned (B) ground otoliths in a transverse plane from a 653 mm spotted gar, noting orientation (P = posterior, A = anterior, V = ventral, D = dorsal). The dashed line represents the portion of the otolith that was removed starting from the anterior (A) side. The nucleus (N) is located in the dorsal region of the otolith, with annuli being estimated in the ventral (V) portion of the otolith.

where otoliths were ground in a plane transverse to the nucleus (Figure 2) using a rotary tool fixed with a grinding bit (#85422, Dremel, Racine WI). The rotary tool was attached to a table, and forceps coated in Tool Dip (Plasti Dip Internation, Blaine MN) were used to securely hold the posterior portion of the otolith during the grinding process. Sagittal otoliths were ground by placing the anterior portion of the otolith on the grinding bit and removing material until the nucleus was even with the apex of the ventral portion of the otolith (Figure 2). Otoliths were then polished using wetted 1600 grit sand paper.

Two readers independently examined the otoliths to estimate ages (Hoff et al. 1997) using an optic-mount digital camera attached to an Olympus dissection microscope (Olympus Corporation, Lake Success NY) and displayed on a high-resolution monitor. Otoliths were selected at random with the readers having no reference to fish length, weight, or sex to reduce bias. Otoliths were then placed in clay and submerged in water to reduce glare. A fiber optic filament with an external light source was used to illuminate annuli. Annuli were first counted

from the nucleus margin to the outer edge, and then repeated to verify the first count. If needed, otoliths were polished multiple times to increase the clarity of and to clearly interpret the outermost annuli (Buckmeier et al. 2012). After otoliths were estimated independently, readers estimated and agreed on ages for both browned and not browned sagittal otoliths. Independently aged otoliths were compared to agreed ages for both browned and not browned sagittal otoliths.

Statistical analyses.—Browned and not browned otoliths were compared using age bias plots to determine if precision was higher using either technique (Campana et al. 1995). Percent agreement (PA) and coefficient of variation (CV) between readers was calculated for both otolith types (Campana et al. 1995). A length frequency histogram was created to visualize the length distribution of Spotted Gar. A von Bertalanffy growth equation (Beverton 1994) was used to model back-calculated total length-at-age using non-linear regression. Non-linear regression was used to describe the weight-length relationship of the population. Due to sample size being low, data for both male and female Spotted Gar were pooled.

Results and Discussion

We collected 24 female and 66 male Spotted Gars, ranging in size between 348 – 846 mm TL (average = 601.1 ± 82.5 mm), from the Lake Thunderbird reservoir (Table 1). Sex ratios were not 1:1 across all ages sampled. This is likely a relic of sampling time and methodology, as gars were targeted during the spawning season. Spawning aggregations of gars typically consist of a greater ratio of males than females (Echelle and Grande 2014; Frenette and Snow 2016). The length-frequency distribution of Spotted Gar appeared to be bimodal, with fewer large (730 - 760 mm TL) individuals than smaller (530 - 630 mm TL) individuals (Figure 3). Gars are known to be sexually dimorphic in body size, with females being larger, on average, than males (Love 2002; McGrath and Hilton 2012; McDonald et al. 2013). In our sample from Lake Thunderbird, female Spotted Gars were, on average, larger (651.7 ± 124.2 mm TL) than males (582.7 ± 48.9 mm TL). The modelled von Bertalanffy growth curve indicates that Spotted Gar from Lake Thunderbird approach maximum length quickly ($K = 0.81$), with individuals in the population growing to half of their expected TL in the first year of life and approaching infinity ($L_{\infty} = 609$ mm) by age 4 (Figure 4). The weight-length relationship suggests that Spotted Gar experience positive isometric growth (Figure 5). It is worth noting that, since sexes were pooled to calculate growth, any sex-specific differences in growth are masked. However, because the sample size of females was low, the difference in back-calculated growth was minimal

compared to when females were excluded from the analysis. Nonetheless, the results hold with the knowledge that gars, including Spotted Gar, grow rapidly during the first year of life (Matthews et al. 2012; David et al. 2015).

Ages assigned using both techniques (not browned and browned) ranged from 1 to 14 years (Table 1). Exact agreements for browned otoliths were 55.4% for reader 1 and 32.1% for reader 2, whereas PA within 1 year was 84.7% for reader 1 and 63.3% for reader 2. CV estimates for browned otoliths were 6.5% for reader 1 and 11.1% for reader 2. Results for not browned otoliths were similar, with exact agreements of 57.1% for reader 1 and 30.7% for reader 2. PA within 1 year for not browned otoliths was 81.3% for reader 1 and 65.1% for reader 2, and CV estimates were 6.1% for reader 1 and 10.6% for reader 2. Age bias plots for both techniques provided similar age estimates and did not consistently under- or overestimate ages (Figure 6), however readers felt that annuli were more clearly visible in browned otoliths. Caution should be used, however, when estimating ages using browned otoliths of Spotted Gar, as false annuli are more defined using this technique and, without experience, are harder to eliminate during estimations.

This study represents the first report of age and size structure of Spotted Gar from the Lake Thunderbird reservoir, and contributes to our understanding of the basic biology of this species in Oklahoma.

Table 1.—Average (± 1 SD) and range (minimum - maximum) of morphometric measurements and age for Spotted Gar from Lake Thunderbird. Measurements are reported for males, females, and for pooled sexes. TL = total length; HL = head length; SnL = snout length; W = weight; A = age.

	Pooled		Female		Male	
	Mean	Range	Mean	Range	Mean	Range
TL (mm)	601.1 ± 82.5	348 - 846	651.7 ± 124.2	356 - 846	582.7 ± 48.9	348 - 674
HL (mm)	143.4 ± 27.5	85 - 240	164 ± 41.7	85 - 240	135.9 ± 13.7	88 - 162
SnL (mm)	96.7 ± 21.2	60 - 173	112.5 ± 30.8	60 - 173	91 ± 11.9	62 - 121
W (kg)	0.82 ± 0.4	0.11 - 2.45	1.14 ± 0.6	0.11 - 2.45	0.70 ± 0.19	0.11 - 1.32
A (yr)	7.1 ± 2.1	1 - 14	6.8 ± 1.8	1 - 9	7.2 ± 2.2	2 - 14




Figure 3.—Length-frequency (TL in mm) histogram for Spotted Gar captured from Lake Thunderbird during spring of 2014.




Figure 4.—Length and age for Spotted Gar collected from Lake Thunderbird during spring of 2014. The solid line represents the modeled von Bertalanffy growth curve. Error bars represent SE.




Figure 5.—Length-weight relationship for Spotted Gar captured from Lake Thunderbird during spring of 2014. The solid line represents the fit between L-W.




Figure 6.—Age bias plots between readers for (A) browned and (B) not browned sagittal otoliths of Spotted Gar from the Lake Thunderbird reservoir. The dashed line represents a 1:1 relationship between readers 1 and 2. Bias between readers is represented by the solid line.

Acknowledgments

The authors thank those individuals that assisted with collection of specimens, lab work, and forage collection including Clayton Porter, Donnie King, Jeff Tibbets, Amie Robison, and Michael Hollie; Kurt Kuklinski (ODWC) for reviewing an early draft of this manuscript; the University of Oklahoma's ARF steering committee for allowing the use of the holding ponds; and Edie-Marsh-Matthews for providing direction during the project. The Oklahoma Department of Wildlife Conservation provided financial support.

This project was approved by IACUC no. R13-003 from the University of Oklahoma.

References

- Beverton RJH. 1994. Notes on the use of theoretical models in the study of dynamics of exploited fish populations. United States Fishery Laboratory, Beaufort (NC), Miscellaneous Contribution (2), 159 pp.
- Buckmeier DL, Smith NG, Reeves KS. 2012. Utility of alligator gar age estimates from otoliths, pectoral fin rays, and scales. *T Am Fish Soc* 141(6):1510-1519.
- Campana SE, Annand MC, McMillan JL. 1995. Graphical and statistical methods for determining the consistency of age determinations. *T Am Fish Soc* 124(1):131-138.
- David SR, Kik IV RS, Diana JS, Rutherford ES, Wiley MJ. 2015. Evidence of countergradient variation in growth of Spotted Gars from core and peripheral populations. *T Am Fish Soc* 144(4):837-850.
- Echelle AA, Riggs CD. 1972. Aspects of the early life history of Gars (*Lepisosteus*) in Lake Texoma. *T Am Fish Soc* 101(1):106-112.
- Echelle AA, Grande L. 2014. *Lepisosteidae: Gars*. In: Warren ML, Burr BM, editors. *Freshwater Fishes of North America*. John Hopkins University Press, Baltimore. p 243-278.
- Ferrara AM, Irwin ER. 2001. A standardized procedure for internal sex identification in *Lepisosteidae*. *N Am J Fish Manage* 21(4):956-961.
- Franssen NR, Tobler M. 2013. Upstream effects of a reservoir on fish assemblages 45 years following impoundment. *J Fish Biol* 82(5):1659-1670.
- Frenette BD, Snow R. 2016. Natural habitat conditions in a captive environment lead to spawning of Spotted Gar. *T Am Fish Soc* 145(4):835-838.
- Glass WR, Corkum LD, Mandrak NE. 2011. Pectoral fin ray aging: an evaluation of a non-lethal method for aging and its application to a population of the threatened Spotted Gar. *Environ Biol Fish* 90(3):235-242.

- Hoff GR, Logen DJ, Douglas MF. 1997. Otolith morphology and increment validation in young Lost River and Shortnose Suckers. *T Am Fish Soc* 126(3):488-494.
- Long JM, Snow RA. 2016. Ontogenetic development of otoliths in Alligator Gar (*Atractosteus spatula*). *T Am Fish Soc* 145(3): 537-544.
- Love JW. 2002. Sexual dimorphism in Spotted Gar *Lepisosteus oculatus* from southeastern Louisiana. *Am Midl Nat* 147(2):393-399.
- Matthews WJ, Shelton WL, Marsh-Matthews E. 2012. First-year growth of Longnose Gar (*Lepisosteus osseus*) from zygote to autumn juvenile. *Southwest Nat* 57(3):335-337.
- McDonald DL, Anderson JD, Hurley C, Bumguardner BW, Robertson CR. 2013. Sexual dimorphism in Alligator Gar. *N Am J Fish Manage* 33(4):811-816.
- McGrath PE, Hilton EJ. 2012. Sexual dimorphism in Longnose Gar *Lepisosteus osseus*. *J Fish Biol* 80(2):335-345.
- NatureServe. 2016. NatureServe Web Service [online]. Available from <http://services.natureserve.org>. (Accessed: August 5, 2016).
- Scarnecchia DL. 1992. A reappraisal of Gars and Bowfin in fishery management. *Fisheries* 17:6-12.
- Secor DH, Dean JM, Laban EH. 1992. Otolith removal and preparation for microstructural examination. In: Stevenson DK, Campana SE, editors. Otolith microstructure examination and analysis. Canadian Special Publication of Fisheries and Aquatic Sciences 117. p 19-57.
- Staton SK, Boyko AL, Dunn SE, Burrige M. 2012. Recovery strategy for the Spotted Gar (*Lepisosteus oculatus*) in Canada (Proposed). Fisheries and Oceans Canada, Species at Risk Act Recovery Strategy Series, Ottawa.
- Tyler JD, Granger MN. 1984. Notes on food habits, size, and spawning behavior of Spotted Gar in Lake Lawtonka, Oklahoma. *Proc Okla Acad Sci* 64:8-10.

Received August 14, 2016 Accepted November 21, 2016