

VASCULAR PLANTS OF SOUTHEASTERN OKLAHOMA FROM THE SANS BOIS TO THE KIAMICHI MOUNTAINS

Submitted to the Faculty of the Graduate College of the Oklahoma State University
in partial fulfillment of the requirements for the Degree of Doctor of Philosophy
May 1969

Francis Hobart Means, Jr.
Midwest City, Oklahoma
Current Email Address: fhmeans@cox.net

The author grew up in the prairie region of Kay County where he learned to appreciate proper management of the soil and the native grass flora. After graduation from college, he moved to Eastern Oklahoma State College where he took a position as Instructor in Botany and Agronomy. In the course of conducting botany field trips and working with local residents on their plant problems, the author became increasingly interested in the flora of that area and of the State of Oklahoma. This led to an extensive study of the northern portion of the Ouachita Highlands with collections currently numbering approximately 4,200. The specimens have been processed according to standard herbarium procedures. The first set has been placed in the Herbarium of Oklahoma State University with the second set going to Eastern Oklahoma State College at Wilburton.

Editor's note: The original species list included habitat characteristics and collection notes. These are omitted here but are available in the dissertation housed at the Edmon-Low Library at OSU or in digital form by request to the editor. [SS]

PHYSICAL FEATURES

Location and Area

The area studied is located primarily in the Ouachita Highlands of eastern Oklahoma. The specific area is generally bounded on the west by State Highway 2, on the south by the Kiamichi Mountains, on the east by the Oklahoma-Arkansas State Line, and on the north by the Sans Bois Mountains. The area includes the southern two-thirds of Latimer County, the southern half of LeFlore County, and the northeast corner of Pushmataha County.

Most of the area is mountainous with prairie sites lying generally west to east in narrow valleys. One large prairie site lies in northern Latimer County and central LeFlore County, between the Sans Bois and

Winding Stair Mountain ranges. A second large valley lies across the southern part of Latimer and LeFlore counties between the Winding Stair and Kiamichi mountain ranges.

Geology

The Sans Bois Mountains of northern Latimer and LeFlore counties are primarily Savanna and McAlester formations of the Krebs group, dating from the Pennsylvanian (Snider 1917). The valleys to the west and east of Wilburton in Latimer County and extending eastward into LeFlore County are alluvium and low terrace deposits underlain by Pennsylvanian strata (Snider 1917). These valleys are generally associated with Gaines Creek in Latimer County, Fourche Maline Creek in Latimer and LeFlore

counties, and Poteau River in LeFlore County. These also are associated with the coal basin of eastern Oklahoma, with numerous coal deposits occurring in the Pennsylvanian strata.

The Ouachita Overthrust, forming the Winding Stair Mountain Range, is of Cambrian to Lower Pennsylvanian rocks formed during the second period of mountain formation in eastern Oklahoma (Dott 1928). The leading edge, to the north and west, formed a great arc and is now known as the Choctaw Fault (Dott 1928), which runs from west to east across central Latimer and LeFlore counties. The Wapanucka limestone outcrops along this fault through Latimer County and dates to the early Pennsylvanian (Fellows 1964). To the south of the Choctaw Fault lie the Winding Stair Mountains. They are heavily faulted, with alternating layers of sandstones and shales. The principal strata are the Atoka Formation of Pennsylvanian, the Jackfork of Upper Mississippian, the John's Valley Formation of Upper Mississippian and Lower Pennsylvanian, and the Stanley Shale of Upper Mississippian (Fellows 1964). Characteristically the valleys are formed from shales and the mountains from sandstones. The deeper riverbeds contain belts of Pleistocene materials. The Atoka Formation is principally gray shale with sandstones distributed throughout, while the Jackfork is composed of heavy, massive beds of brown sandstone separated by thinner bands of gray shale (Snider 1917). The Atoka strata are more resistant to erosion than is the Stanley Shale. The John's Valley Formation is gray-green clay shale with interbedded sandstone that is easily eroded and is a valley former (Snider 1917). The Stanley Shale is a bluish, greenish-black slaty shale with thin sandstone layers and considerable chert (Snider 1917). Some Caney Shale, which is black and green in color, is also present.

The Winding Stair Mountains are separated from the Kiamichi Mountains by

a valley through which flows the Kiamichi River. This valley runs from Big Cedar in LeFlore County westward through Talihina, across Latimer County, and into Pushmataha County near Clayton. It is formed from the Stanley Shale of Upper Mississippian (Snider 1917) that is easily eroded and forms valley floors.

Lying to the west of Talihina in southeast Latimer County and along the northeastern edge of Pushmataha County are the Potato Hills. They are remnants of Ordovician black shales and sandstones underlain by shales of Lower Pennsylvanian (Snider 1917). At many places the Arkansas Novaculite-Stanley complex is present. The Ordovician consists of bluish, greenish, and white chert with thin cherty and slaty shales with thin lenticels of limestone (Snider 1917). The Kiamichi Mountains are Jackfork Shale of Pennsylvanian underlain by Stanley Shale (Snider 1917). The sandstone is resistant to weathering. The faulting is severe, and these mountains are more rugged than the Winding Stair Mountains. Rich Mountain, of southeast LeFlore County, is of like composition. It is the highest in the area, reaching an elevation of approximately 3000 feet (914.4 m) above sea level (Snider 1917), which is approximately 2000 feet (609.6 m) above the streams at its base. The lower elevations in the area are approximately 500 feet (152.4 m) above sea level.

Topography by Counties

According to Snider (1917), Latimer County covers approximately 735 sq. miles (1903.64 sq. km), lying in an area of Pennsylvanian rocks with the southern part in the Ouachita Mountain region. Both the northern and southern parts consist of alternating sandstones and shales of considerable thickness, folded into steep, northeast-southwest folds. The southern formations are the oldest and are steeper.

The northern part is drained by Fourche Maline eastward into the Poteau River; the southern part drains into the Kiamichi River through several small tributaries; and the southwestern part drains northwest into the Canadian River by way of Gaines Creek (Figure).

LeFlore County is one of the larger counties in the state, covering approximately 1614 sq. miles (4180.24 sq. km). The northern part lies in the Arkansas Valley geologic and physiographic province while the southern part lies in the Ouachita Mountains. The formations are the same as those of Latimer County, with the Ouachita Mountains being especially rough. Included are the Winding Stair, Kiamichi, and Jackfork mountains. The middle and northern part drains into the Arkansas River by way of the Poteau River and its tributaries. The southern part drains into the Kiamichi River, which lies north of Kiamichi Mountain. The southern edge, south of Kiamichi Mountain, is drained by Little River toward the Red River.

Pushmataha County covers approximately 1430 sq. miles (3703.68 sq. km.). It lies in the Ouachita Mountains except for the southwest corner. The hills are rugged with much surface sandstone, separated by narrow valleys. The Jackfork Mountains and the Potato Hills are the areas of highest elevation in the county. The eastern part of the county drains toward the Red River by means of Little River. Most of the county is drained toward the Red River by Kiamichi River and its tributaries. The Kiamichi has its origin in Arkansas, flows westward, then southward, then southeast to the Red River. The Kiamichi River is paralleled by Little River to the east, which has its origin in the southwest corner of LeFlore County.

Soils

According to Gray and Galloway (1959) there are four principal soil series within the

Ouachita Highlands, with several additional localized series. They are acid red-yellow podzolic soils developed from gray and brown shales and sandstones. The surface soils are generally light colored and strongly leached. The major soil series are the Hector-Pottsville, Enders-Conway-Hector, Atkins-Pope, and the Parsons-Dennis-Bates.

South of the Choctaw Fault, much of the mountains are rough with some of the formations steeply tilted (as much as 60 degrees to the horizontal). On these mountains, forest vegetation can easily penetrate the more weathered layers of the slopes. The Sans Bois Mountains have soils of the same series as the Winding Stair and Kiamichi mountains, but the strata are more horizontal, which results in poorer forest sites.

The soil association characteristic of the Sans Bois, Winding Stair, and Kiamichi mountain ranges is the Hector-Pottsville series (see Fig.). In the Hector series, the topsoil is a dark brown sandy loam, characteristically formed on hills and mountains under forest type vegetation. In the Pottsville series, topsoil is a brown, fine sandy loam or loam a few inches deep, formed on hills and mountains under forest vegetation. In both Hector and Pottsville, it is not uncommon for slopes to be steep. Ledges and surface rocks are also a common occurrence.

The soils of the Potato Hills include the Clebit series in addition to the Hector and Pottsville series. The Clebit topsoil varies from a dark gray-brown stony silt loam to a pale brown silt loam, formed under forest vegetation of rocky steep slopes.

The soil of the prairie west of Wilburton, in Latimer County, is the Parson-Dennis-Bates Association (see Figure). This association also forms part of the prairie north of the Kiamichi River, south and east of Talihina. The Parson topsoil is a grayish brown medium acid silt loam. The topography is nearly level to gentle slopes and is covered by tall grass

vegetation. The permeability of the soil is slow; it is seasonally wet and of low fertility. The Dennis series topsoil is dark grayish brown, medium acid silt loam of low fertility, and is subject to erosion. The Bates series topsoil is a dark grayish brown loam or fine sandy loam of medium acidity. Shallow spots with surface rock are common and the soil is easily eroded.

The large prairie of eastern Latimer County and northern LeFlore County, north of the Ouachita Highlands, is of the Enders-Conway-Hector Association (see Figure). The Enders soil is a brown, fine, sandy loam found on gentle slopes and ridges and is of low fertility, erosive, and droughty. The Conway series is a brown silt loam. Parent material is gray and brown clay and sandy shales of gentle slopes and valleys. It is of low fertility, slow draining, and it commonly has silt mounds. The Hector series, which is a part of this association, has been previously discussed.

The Poteau River valley has soils of the Atkins-Pope Association (see Figure). The Atkins series is a light gray acid silt loam or gray loam mottled with brown and yellow. The parent materials are gray acid mottled clay loams, loams, and loamy alluviums. The terrain is flood plain, subject to overflow, and is of low fertility with poor drainage. The Pope soil is a brown acid fine sandy loam. Parent materials are brown stratified alluviums of flood plains and naturally elevated dikes. It is subject to overflow, is of low fertility, but is sandier and better drained than the Atkins.

CLIMATE

The climate of the Ouachita Highlands is of the continental type. It is moderated by seasonal influences of warm moist winds from the Gulf of Mexico. The annual temperature extremes range from a few degrees below zero to 103° F (-20° to 40° C).




Figure Generalized soil map of southeastern Oklahoma. Symbols are: PDB – Parson, Dennis, Bates; ECH – Enders, Conway, Hector; HP – Hector, Pottsville; and AP – Adkins, Pope (Gray and Galloway 1959).

The average temperature for the years 1962 through 1967 was approximately 62° F (17° C; Table I).

Precipitation is high, ranging from just under 40 inches (101.6 cm) in the northern part to nearly 50 (127 cm) inches in the southern mountainous part (Table II). During the relatively dry year of 1963, just over 20 inches (50.8 cm) fell at Wilburton, whereas in 1967, over 62 inches (157.5 cm) fell at the Kiamichi Tower on Kiamichi Mountain in Southwest LeFlore County (Table II).

The distribution of rainfall is more uniform over the entire year than is usual for the rest of Oklahoma. Spring is characterized by heavy rainfall, resulting in considerable local flooding. The summer months often become droughty with

moisture again being plentiful during the fall. Eastern Oklahoma and the adjoining states receive, on the average, more precipitation in the spring than any other state east of the Rocky Mountains (Wahlgren 1941).

High summer temperatures usually occur with clear skies and are accompanied

by light wind. In winter, occasional sleet, ice storms, or snows occur but are few in number and of short duration. The last killing frost in spring falls in March to late April, with the first frost in fall occurring in late October.

TABLE I
AVERAGE TEMPERATURES

Station	1962	1963	1964	1965	1966	1967
Clayton 2N			63.3M			
Kiamichi Tower						
Poteau	62.3	63.1	62.5	62.0	60.9	62.1
Smithville 2NNW	60.5	60.9				60.1M
Wilburton	62.2	63.4	62.5	62.4M	59.7M	
Wister Dam	63.1	63.7	63.3	63.8M	60.4M	60.7

Source: U.S. Dept. of Commerce, Weather Bureau, Climatological Data, Oklahoma, Annual Summaries, 1962-1967. Temperature averages followed by an M indicate one or more months of missing data.

TABLE II
AVERAGE PRECIPITATION

Station	1962	1963	1964	1965	1966	1967
Clayton 2N	48.5	29.4E	50.4	36.0E	37.0E	53.8E
Kiamichi Tower	59.0E	28.0E	45.4	58.2E	44.1E	62.1
Poteau	41.8	20.9	38.6	31.6	33.1	49.6
Smithville 2NNW	51.3	29.7	43.5	56.9	39.2E	59.2
Wilburton	45.7	20.7	41.9	35.7	45.9	48.8E
Wister Dam	38.7	22.0	40.3	32.6	36.9	50.5

Source: U.S. Dept. of Commerce, Weather Bureau, Climatological Data, Oklahoma, Annual Summaries, 1962-1967. Precipitation averages followed by an E indicate one or more months of missing data.

TAXONOMIC HISTORY

The first plant collecting in the Ouachita Highlands of eastern Oklahoma was done by Thomas Nuttall in 1819. During a stay of several weeks at Ft. Smith on the Arkansas River in western Arkansas, he made several short collecting trips into surrounding territory. On May 16, 1819, Nuttall left Ft. Smith with Major Bradford and a company of soldiers on a trip to the confluence of the Kiamichi River and the Red River. They followed the Poteau and Kiamichi rivers, crossing the mountains that separate the two drainage systems.

The following are excerpts from *Nuttall's Journal of his Travels into the Arkansas Territory* (as reprinted in *Early Western Travels*, Volume XIII. Thwaites 1905).

[April] 27. Yesterday I took a walk of about five miles up the banks of the Pottoe [Poteau], and found my labour well repayed by the discovery of several new or undescribed plants... The whole expanse of forest, hill and dale was now richly enameled with a profusion of beautiful and curious flowers; among the most conspicuous was the charming Daisy of America [*Astranthium integrifolium* (Michx.) Nutt.] of a delicate lilac colour, and altogether corresponding in general aspect with the European species; intermingled, appears a new species of Phlox, the *Verbena subletia*, and the esculent *Scilla [camassia]*.

Nuttall made interesting notes about the terrain and flora. After passing the Poteau River, he noted the conic shape of Sugar Loaf Mountain and Cavanah Mountain, likening them to the Allegheny Mountains (Thwaites 1905). On May 17, he recorded the following:

These vast plains, beautiful almost as the fancied Elysium, were now enamelled with innumerable flowers, among the most splendid of

which were the azur Larkspur [*Delphinium carolinianum* Walt.], gilded Coreopsides [*Coreopsis*], Rudbeckias [*Rudbeckia*], fragrant Phloxes, and the purple *Psilotria*.

After crossing the divide from the Poteau River to the Kiamichi River, he again likened the ridges to the Allegheny of Pennsylvania, noting that they were rocky and thinly wooded with pines and oaks (Thwaites 1905).

On his return trip to Ft. Smith, Nuttall notes passing with great difficulty along the summit of a mountain covered with thickets of "dwarf oaks (*Quercus chinquapin*, *Q. montana* and *Q. alba*), none of them scarcely exceeding the height of a man" (Thwaites 1905).

Many other botanists traveled and collected plants in Oklahoma. Zina Pitcher, a surgeon in the U. S. Army, apparently traveled the same general route as did Nuttall (McKelvey 1955). Melines C. Leavenworth, Heinrich Karl Beyrich, Charles Joseph Latrobe, and Edward James all collected in Oklahoma but passed by the Ouachita Highlands while enroute to more western or southern destinations. (McKelvey 1955).

G. D. Butler collected *Isoetes* at Limestone Gap, approximately 70 miles north of Texas and 100 miles west of Arkansas (Butler 1878). This is probably the present town of Gap in northern Atoka County, situated in a break of the Ouachita Highlands.

Stevens collected plants from the vicinity of Page, Oklahoma, in southeastern Oklahoma, during 1913 prior to his and Shannon's joint publication of plant life in Oklahoma (Stevens and Shannon 1917).

Palmer (1924) made a study of the ligneous flora of Rich Mountain in southeastern Oklahoma and Featherly (1928) listed the grasses of Oklahoma. In addition, occasional collections from the area of study by E. Little, R. Stratton, and

G. Goodman are in the herbarium of Oklahoma State University. Collections of significance include those of U. T. Waterfall.

ECOLOGICAL CONSIDERATIONS

The vegetation of the Ouachita Highlands is the Oak-Hickory Association of the Deciduous Forest Formation (Bruner 1931). This association is composed of two communities that are similar and intergrade considerably. One community is the upland forests of the rough hills and mountainous areas and the other is the lowland forest of stream valleys and more mesic lower slopes. Overlapping into the forests, primarily in the valleys, is the tall grass prairie. There is intergradation to a limited extent between the lowland forests and the prairie community.

The oak-hickory forest is most extensive on the lower slopes and level fertile valleys. The dominant species are *Quercus shumardii*, var. *schneckii*, *Q. nigra*, *Q. falcata*, var. *falcata*, *Q. velutina*, *Q. stellata*, *Carya aquatica*, *C. cordiformis*, *C. myristicaeformis*, and *C. texana*. Other species commonly present in the valleys and lower slopes include *Acer saccharum*, *A. saccharinum*, *A. negundo*, *Diospyros virginiana*, *Sassafras albidum*, *Liquidambar styraciflua*, *Juglans nigra*, *Prunus serotina*, *Robinia pseudoacacia*, *Nyssa sylvatica*, *Ostrya virginiana*, *Tilia neglecta*, *Quercus macrocarpa*, *Q. muhlenbergii*, *Q. lyrata*, *Q. alba*, *Ulmus alata*, *Carya illinoensis*, *Celtis laevigata*, *Pinus echinata*, *Salix caroliniana*, *S. nigra*, *Platanus occidentalis*, *Gleditsia tricanthos*, *Fraxinus americana*, *Maclura pomifera*, and *Salix interior*.

Characteristic dominant species of the more xeric upland sites include *Quercus velutina*, *Q. stellata*, *Q. palustris*, *Q. marilandica*, *Carya cordiformis*, *C. texana*, *Ulmus alata*, and *Pinus echinata*. Other species, including shrubs, present in the upland forest include the following: *Aesculus glabra*, *Ascyrum hypericoides*, *Vaccinium arboretum*, *V. stamineum*, *Bumelia lanuginosae*, *Ceanothus herbaceus*,

Hypericum spathulatum, *H. punctatum*, *Ilex deciduas*, and *Rhus copallina*. Shrubs more characteristic of the more mesic lower slopes include *Crataegus crus-galli*, *C. spathulata*, *Cornus drummondii*, *C. obliquae*, *Prunus mexicana*, *P. americana*, *Rhus toxicodendron*, *R. glabra*, *R. radicans*, and *Virburnum prunifolium*.

Shrubs of the mesic lower slopes and valleys include *Cornus florida*, *Alnus serrulata*, *Betula nigra*, *Ostrya virginiana*, *Carpinus caroliniana*, *Callicarpa americana*, *Hamamelis vernalis*, *Prunus serotina*, *Amorpha fruticosa*, *Hydrangea arborescens*, and *Asimina triloba*. *Magnolia acuminata* is found only on the mesic northern slope of Rich Mountain in Southeast LeFlore County. *Ilex opaca* is restricted to wet sandy loam soils along Kiamichi River and Little River.

Common lianas found in the Oak-Hickory Association include the following: *Clematis versicolor*, *Calyocarpum lyoni*, *Cocculus carolinus*, *Vitis rotundifolia*, *V. vulpina*, *V. acerifolia*, *V. aestivalis*, *Parthenocissus quinquefolia*, *Menisperma canadense*, *Ampelopsis cordata*, *Berberis scandens*, *Cissus incisa*, *Smilax glauca*, *S. bona-nox*, *S. rotundifolia*, *Rhododendron oblongifolium*, and *Campsis radicans*.

The herbaceous flora varies with the seasons and the density of the forest. The prevernal and vernal species include the following: *Sanguinaria canadensis*, *Podophyllum peltatum*, *Polygonatum canaliculatum*, *Arisaema dracontium*, *Erythronium americanum*, *Danthonia spicata*, *Panicum sphaerocarpon*, *Carex brevior*, *C. caroliniana*, *C. lurida*, *Trillium viride*, *Viola pedata*, var. *lineariloba*, *V. sororia*, *V. kitaibeliana*, *Valerianella longiflora*, *V. stenocarpa*, *Ranunculus hispidus*, *R. fascicularis*, *Antennaria plantaginifolia*, *Senecio obovatus*, *Oenothera laciniata*, *Anemonella thalictroides*, *Lepidium virginianum*, and *Callirhoe alcaeoides*.

Common estival species include the following: *Silene stellata*, *Salvia lyrata*, *Monarda fistulosa*, *Teucrium canadense*, *Geum canadense*, *Tovara virginiana*, *Boehmeria cylindrica*, *Utriclea chamaedryoides*, *Commelina communis*, *Polygonum hydropiperoides*, var. *opelousanum*, *Rumex crispus*,

R. pulcher, *Froelichia gracilis*, *Tephrosia virginiana*, *Zizia aurea*, *Cassia fasciculata*, *Clitoria mariana*, *Desmodium sessilifolium*, *Elymus canadensis*, *Panicum hians*, *Eleocharis obtusa*, *Lobelia spicata*, and *Passiflora incarnata*.

The serotinal species include the following: *Lobelia cardinalis*, *Aster azureus*, *Boltonia diffusa*, *Elephantopus carolinianus*, *Eupatorium coelastinum*, *E. serotinum*, *Iresine rhizomota*, *Impatiens capensis*, *Agastache nepetoides*, *Helianthus hirsutus*, *Uniola latifolia*, *Tridens flavus*, *Croton monanthogynus*, *Euphorbia corollata*, *E. supine*, *Pycnanthemum albescens*, *Prunella vulgaris*, *Plantago rugelii*, *Coreopsis grandiflora*, and *Solidago delicatula*.

Subclimax prairie is found between the Winding Stair Mountains and the Sans Bois Mountains, and between the Kiamichi Mountains and the Winding Stair Mountains.

Dominant prairie species are as follows: *Andropogon gerardi*, *A. scoparius*, *Sorghastrum nutans*, and *Panicum virgatum*. Other common species include *Andropogon saccharoides*, *A. ternarius*, *Setaria geniculata*, *Echinochloa crusgalli*, *Panicum anceps*, *P. agrostoides*, var. *condensatum*, *Paspalum setaceum*, *Agrostis hyemalis*, *Aristida oligantha*, *Spehnopholis obtusata*, *Tridens strictus*, *Carex amphibola*, var. *turgid*, *C. lupuliformis*, *Scirpus lineatus*, *Aristida longespica*, *Elymus virginicus*, *Manisuris cylindrical*, *Eragrostis trichodes*, *Bromus secalinus*, *Festuca octaflora*, and *Hordeum pussillum*.

Prevernal and vernal species of the prairies include the following: *Sisyrinchium campestre*, *Hypoxis hirsuta*, *Tradescantia obiensis*, *T. ernestiana*, *Baptisia leucophaea*, *B. nuttallina*, *B. sphaerocarpa*, *Collinsia violacea*, *Ranunculus fascicularis*, *Bromus japonicus*, *B. mollis*, *Penstemon arkansanus*, *P. digitalis*, *Claytonia virginica*, *Anemone caroliniana*, *Linaria canadensis*, *Camassia angusta*, and *Luzula bulbosa*.

Species that are a little later but still vernal include *Daucus pusillus*, *Ptilimnium nuttallii*, *Potentilla canadensis*, *Amsonia tabernaemontana*, *Phacelia hirsuta*, *Astranthium integrifolium*, *Phlox pilosa*, *Psoralea psoralioides*,

P. tenuiflora, *Silene stellata*, *Astragalus distortus*, *Rosa carolina*, *Stylosanthes biflora*, *Polygala incarnate*, *Acalypha virginica*, and *Verbena canadensis*.

Estival species of the prairie include *Rudbeckia grandiflora*, *R. triloba*, *Cicuta maculata*, *Asclepias tuberosa*, *Oenothera laciniata*, *Zizia aurea*, *Eryngium yuccifolium*, *Gaura filiformis*, *Liatris pycnostachya*, *Spiranthes vernalis*, and *Cuscuta cuspidata*.

Some serotinal species of the prairie include *Solidago radula*, *S. rigida*, *Helianthus mollis*, *Vernonia baldwinii*, *Silphium laciniatum*, and *Euphorbia nutans*.

Vegetation associated with the streams, ponds and lakes of the area varies from free floating aquatics to those growing along the edge of water. Common free floating or bottom rooted species include *Nuphar advena*, var. *advena*, *Lemna valdiviana*, *Myriophyllum heterophyllum*, *Utricularia biflora*, *Potamogeton diversifolius*, and *Najas guadalupensis*. Species rooted at the edge of the water include *Typha latifolia*, *Sagittaria gramineum*, *S. ambigua*, *Zizaniopsis miliacea*, *Hydrolea ovata*, *Justicia americana*, *Polygonum pennsylvanicum*, *P. persicaria*, *Ludwigia palustris*, *Eleocharis obtusa*, *E. quadrangulata*, *Rhynchospora corniculata*, and various species of *Carex*.

RANGE EXTENSIONS AND SPECIES OF SPECIAL INTEREST

This chapter covers range extensions and species having a rather restricted distribution in the area studied.

Range Extensions

Bidens aristosa (Michx.) Britt. var. *mutica* (Gray) Gattinger, reported by Waterfall (1954a) for McCurtain County, was collected in early October (Means 2837) near Lake Nahih Wayia in Pushmataha County and (Means 2210) in the Poteau River valley 5 mi. (8.05 km) south of Poteau in LeFlore County.

Carex lactebracteata Waterfall, a new species described by Waterfall (1954a) with the TYPE (Waterfall 11380) from a rocky wooded ridge 16.4 mi. (26.39 km) north of Broken Bow, in McCurtain County; it was collected May 1968 by the author (3252) on rocky wooded hillsides of Cucumber Creek in LeFlore County, approximately 22 mi. (35.41 km) north of the original collection site.

Species of Special Interest

Pinus teada L., although occurring in large stands in Southeast McCurtain County, was collected (Means 2499) as an occasional tree of mixed hardwood forest of Kiamichi River valley approximately 2 mi. (3.22 km) southeast of Tuskahoma in Pushmataha County in October.

Taxodium distichum (L.) Richard, collected in August, 1965 (Means 2066) and early April, 1966 (Means 2403) in shallow water along the banks of Poteau River, near an old home site in LeFlore County. Reproduction has occurred.

Buchloe dactyloides (Nutt.) Engelm., common further west in the grasslands, collected (Means 1521) on clay site along creek 1.5 mi. (2.41 km) east of the Latimer-LeFlore county line along Highway 270 in LeFlore County in early June, 1965.

Cynosurus echinatus L., naturalized from Europe; collected (Means 2597) in open woods on a rocky hillside near a stream 1 mi. (1.61 km) south of Clayton in Pushmataha County in early June, 1968.

Xyris torta J. E. Smith, var. *occidentalis* Malme, collected (Waterfall 10547) in a srew 1.1 mi. (1.77 km) west of Talihina in Latimer County, October 14, 1951.

Populus deltoides Marsh., although fairly common in central and western Oklahoma, collected (Means 2361) at the base of wooded north slope of a ridge north of Eastern Oklahoma State College at Wilburton in Latimer County, April 5, 1966.

Brasenia schreberi Gmel., abundant in upper end of Lake Nanih Wayia, June 16, 1968 (Means 3608) in Pushmataha County and less commonly in a farm pond 2 mi. (3.22 km) east of Buffalo Valley School, July 13, 1968 (Means 3790) in southeastern Latimer County.

Magnolia acuminata L., reported by Palmer (1924) as occurring on the lower north slopes of Rich Mountain in LeFlore County, collected in June 1932 (Stevens 2771); June 1968 (Means 3553); and May 1968 (Means 3279) only at that site, approximately 0.5 mi. (0.8 km) west of the Arkansas border in early June.

Asimina triloba (L.) Dunal, indicated as common near Page, Oklahoma by Palmer (1924), collected in fruit (Means 790) in the wooded valley of Cucumber Creek in LeFlore County, August 17, 1963.

Drosera annua Reed, collected (Barclay and Doty sin. Num.) May 1961, on sides of low mounds in prairie along Highway 2 approximately 4 mi. (6.44 km) north of Clayton in Pushmataha County.

Sedum nuttallianum Raf., collected May 23, 1966 (Means 2453) from a rather dense stand on a wet weather seep at the edge of a blue shale outcrop along the highway, approximately 1.2 mi. (1.93 km) west and 0.7 mi. (1.13 km) north of Tuskahoma in Pushmataha County.

Ribes cynosbati L., reported by Palmer (1924) from Rich Mountain, collected (Means 2507) from mixed hardwood forest of Kiamichi River valley 0.5 mi. (0.8 km) south and 1.5 mi. (2.41 km) east of Tuskahoma in Pushmataha County, May 22, 1966 and (Means 2893) from the rocky north slope of Kiamichi Mountain 1 mi. (1.61 km) south of Big Cedar in LeFlore County, April 15, 1967.

Andrachne phyllanthoides (Nutt.) Coulter, occasional shrubby plant of rock-strewn small streams, collected (Means 3656) along edge of small rocky stream 1 mi. (1.61 km) northwest of Albion in Pushmataha County, June 30, 1968. Also collected (Waterfall

8542) along rocky stream west of Talihina October 11, 1964 and (Waterfall 17171) along a rocky stream 9 mi. (14.48 km) north of Tuskahoma, August 9, 1948, both in Latimer County.

Ilex opaca Ait., collected (Means 1408) May 26, 1965 and (Means 2436) April 22, 1966 in deep sandy soil of Kiamichi River valley 0.7 mi. (1.13 km) south of Big Cedar; (Clark 350) May 3, 1935 and (Stevens 1406) April 1914 in wooded valley near Page; (Means 3220) in Little River valley 6 mi. (9.65 km) southeast of Nahoba May 27, 1968; (Sellers sin.num.) July 16, 1966 9 mi. (14.48 km) northwest of Clayton in Pushmataha County.

Proserpinaca palustris L., var. *crebra* Fern. & Grisc., collected (Means 3585) in a stream 1

mi. (1.61 km) south of Clayton in Pushmataha County, June 16, 1968.

Myriophyllum pinnatum (Walt.) BSP., collected (Means 3401) in shallow water of Fourche Maline Creek 4 mi. (6.44 km) south of Red Oak in Latimer County, May 28, 1968.

Liatris elegans (Walt.) Willd., collected (Means 3944) August 26, 1968 and (Waterfall 147) July 31, 1932 in native prairie west of Albion in Pushmataha County; (Stratton 604) September 1927 and (Waterfall 15173) October 11, 1958 in prairies east of Wilburton in Latimer County.

Rudbeckia maxima Nutt., collected (Means 1507) from a wet prairie site 1 mi. (1.61 km) west of Red Oak in Latimer County, June 16, 1965.

**TABULAR VIEW OF THE FAMILIES
GENERA (G) and SPECIES AND SUBSPECIFIC TAXA (SS)**

Family	G	SS	Family	G	SS
OSMUNDACEAE	1	1	PHYTOLACCACEAE	1	1
POLYPODIACEAE	11	13	AIZOACEAE	1	1
PINACEAE	3	4	PORTULACACEAE	3	4
TYPHACEAE	1	3	CARYOPHYLLACEAE	6	12
SPARGANIACEAE	1	1	NYMPHACEAE	4	7
ZOSTERACEAE	1	2	RANUNCULACEAE	7	16
NAJADACEAE	1	1	BERBERIDACEAE	1	1
ALISMACEAE	3	6	MENISPERMACEAE	2	2
GRAMINEAE	52	149	MAGNOLIACEAE	1	1
CYPERACEAE	6	59	ANONACEAE	1	1
ARACEAE	1	2	LAURACEAE	2	3
LEMNACEAE	2	2	PAPAVERACEAE	1	1
XYRIDACEAE	1	2	FUMARIACEAE	1	2
COMMELINACEAE	2	13	CRUCIFERAE	11	14
JUNCACEAE	2	15	CAPPARIDACEAE	2	2
LILACEAE	14	25	DROSERACEAE	1	1
AMARYLLIDACEAE	3	4	CRASSULACEAE	1	1
DIOSCOREACEAE	1	2	SAXIFRAGACEAE	6	6
IRIDACEAE	2	3	HAMAMELIDACEAE	2	2
MARANTACEAE	1	1	PLATANACEAE	1	1
ORCHIDACEAE	2	4	ROSACEAE	10	32
SAURURACEAE	1	1	LEGUMINOSAE	25	71
SALICACEAE	2	5	GERANIACEAE	1	1
JUGLANDACEAE	2	8	OXALIDACEAE	1	5
BETULACEAE	4	4	LINACEAE	1	2
FAGACEAE	3	19	ZYGOPHYLLACEAE	1	1
ULMACEAE	2	4	RUTACEAE	1	1
MORACEAE	2	3	MELIACEAE	1	1
UTRICACEAE	4	4	POLYGALACEAE	1	5
LORANTHACEAE	1	1	EUPHORBIACEAE	9	26
ARISTOLACHIACEAE	1	1	CALLITRICHACEAE	1	1
POLYGONACEAE	5	20	ANACARDIACEAE	1	5
CHENOPODIACEAE	2	5	AQUIFOLIACEAE	1	2
AMARANTHACEAE	3	6	CELASTRACEAE	1	2
NYCTAGINACEAE	1	2	STAPHYLEACEAE	1	1

ACERACEAE	1	4
HIPPOCASTANACEAE	1	1
SAPINDACEAE	1	1
BALSAMINACEAE	1	1
RHAMNACEAE	3	4
VITACEAE	4	11
TILIACEAE	1	3
MALVACEAE	5	8
GUTTIFERAE	2	9
CISTACEAE	1	2
VIOLACEAE	1	15
PASSIFLORACEAE	1	3
CACTACEAE	1	2
LYTHRACEAE	3	3
MELASTOMACEAE	1	1
ONAGRACEAE	4	15
HALORAGACEAE	2	4
UMBELLIFERAE	17	21
CORNACEAE	1	3
NYSSACEAE	1	1
ERICACEAE	3	6
PRIMULACEAE	3	3
SAPOTACEAE	1	1
EBENACEAE	1	2
STYRACACEAE	1	1
OLEACEAE	2	3
LOGANIACEAE	3	3
GENTIANACEAE	2	4
APOCYNACEAE	3	4
ASCLEPIADACEAE	2	12
CONVOLVULACEAE	3	12
POLEMONIACEAE	2	3
HYDROPHYLLACEAE	3	5
BORAGINACEAE	5	5
VERBENACEAE	2	7
LABIATAE	18	28
SOLANACEAE	3	15

SCROPHULARIACEAE	15	27
BIGNONIACEAE	2	2
LENTIBULARIACEAE	1	1
ACANTHACEAE	3	6
PHRYMACEAE	1	1
PLANTAGINACEAE	1	5
RUBIACEAE	5	11
CAPRIFOLIACEAE	4	7
VALERIANACEAE	1	4
CUCURBITACEAE	2	2
CAMPANULACEAE	2	8
COMPOSITAE	56	137
<hr/>		
TOTALS	457	1067

Out of a total of 119 families, the ten families with the largest number of species and subspecies are as follows:

GRAMINEAE	52	149
COMPOSITAE	56	137
LEGUMINOSAE	25	71
CYPERACEAE	6	59
ROSACEAE	10	32
LABIATAE	18	28
SCROPHULARIACEAE	15	27
EUPHORBIACEAE	9	26
UMBELLIFERAE	17	21
POLYGONACEAE	5	20
<hr/>		
TOTALS	213	570

SUMMARY

After moving to Eastern Oklahoma State College as Instructor in Botany and becoming aware of the variety in the local flora, the author began an extensive study of the northern Ouachita Highlands. The author has authenticated approximately 4,500 sheets which have been processed according to standard herbarium procedures. The first set has been placed in the Herbarium of Oklahoma State University with duplicates going to Eastern Oklahoma State College at Wilburton, Oklahoma.

Monographs, revisions, and other taxonomic literature from the Oklahoma State University Library and the personal libraries of Dr. U.T. Waterfall and the author were used in the identification of the specimens.

A total of 1067 species and subspecific taxa representing 457 genera and 119 families were identified. The families having the greatest number of species and subordinate taxa were *Gramineae* 149, *Compositae* 137, *Leguminosae* 71, *Cyperaceae* 59, *Rosaceae* 32, *Labiatae* 28, *Scrophularaceae* 27, *Euphorbiaceae* 26, *Umbelliferae* 21, and *Polygonaceae* 20. These 10 families contain 53% of the total species and subordinate taxa.

In 1969, no taxa were reported as new records for the state, although 17 species were listed as range extensions or of special interest due to their limited distribution.

ACKNOWLEDGEMENTS

The author wishes to express his appreciation to each member of his committee for their interest shown and guidance given. He is especially grateful to the chairman, Dr. U.T. Waterfall for his patience, continued interest and the use of his personal library and card index of research literature.

Means, F.H.

LITERATURE CITED

- Allen, Paul and Theodor Just. 1934. Key and synopsis of the American species of the genus *Chenopodium* L. *Am. Midl. Nat.* 30:47-76.
- Bailey, L.H. 1932. The blackberries of North America. *Gentes Herb.* 2(6):397-423.
- Bailey, L.H. 1945. The genus *Rubus* in North America. *Gentes Herb.* 5(9):591-856.
- Barneby, Rupert C. 1964. *Atlas of North American Astragalus*. Mem. N. Y. Bot. Gard. 13. I:1-596, II:597-1188.
- Beetle, Alan Ackerman. 1947. *Scirpus*. *N. Am. Fl.* 18(8):481-504.
- Benson, Lyman. 1948. A treatise of the North American *Ranunculi*. *Am. Midl. Nat.* 40:1-261.
- Boivin, Bernard. 1944. American *Thalictra* and their Old World allies. *Rhod.* 46:335-349, 469-471, 480-483.
- Britton, N.L. and J.N. Rose. 1919-1923. The *Cactaceae*. *Carnegie Inst. Washington*. Vol.1-4.
- Bruner, W.E. 1931. The vegetation of Oklahoma. *Ecological Monographs* I, (2):101-188.
- Butler, G.D. 1878. A list of some of the most interesting species of plants collected in the Indian Territory. *Bot. Gaz.*, 3:65-68, 74-78.
- Butters, F.K. and E.C. Abbe. 1940. The American varieties of *Rorippa islandica*. *Rhod.* 42:25-32.
- Clewell, A.F. 1966. Native North American species of *Lespedeza* (Leguminosae). *Rhod.* 68:359-405.
- Core, Earl L. 1941. The North American species of *Paronychia*. *Am. Midl. Nat.* 26:269-398.
- Correll, Donovan S. 1956. Ferns and fern allies of Texas. *Contr. Texas Research Foundation* 2:1-188.
- Cronquist, Arthur. 1947. Revision of the North American species of *Erigeron*, north of Mexico. *Britt.* 6(2):121-302.
- Dott, Robert H. 1928. Pennsylvanian paleogeography with special reference to South-Central Oklahoma. *Okla. Geol. Surv. Bul.* No. 40:51-68.

- Erickson, Ralph O. 1943. Taxonomy of *Clematis* section *Viorna*. *Ann. Mo. Bot. Gard.* 30:1-30.
- Ernst, W.R. 1963. The genera of Capparidaceae and Moringaceae of the South East United States. *Journ. Arn. Arb.* 44(1):81-95.
- Featherly, H.I. 1946. Manual of the grasses of Oklahoma. *Bul. Okla. A. & M. College* 43(21):1-137.
- Fellows, L.D. 1964. Geology of the western part of the Winding Stair Range, Latimer and LeFlore counties, Oklahoma. *Okla. Geol. Surv. Circ.* 65:1-102.
- Ferguson, A.M. 1901. *Crotons* of the United States. *Rept. Mo. Bot. Gard.* 12:33-74.
- Fernald, M.L. 1922. Notes on *Sparganium*. *Rhod.* 24:26-33.
- _____. 1931. *Potentilla canadensis* and *Potentilla simplex*. *Rhod.* 33:180-191.
- _____. 1932. The linear-leaved North American species of *Potamogeton* section *Axillares*. *Mem. Am. Acad. Arts & Sciences* 17:1-183.
- _____. 1933. Types of some American species of *Elymus*. *Rhod.* 35:187-198.
- _____. 1934. *Draba* in temperate northeastern America. *Rhod.* 36:241-261, 285-305, 314-344, 353-371, 392-404.
- _____. 1938a. VIII New species, varieties and transfers. *Rhod.* 40:334-338.
- _____. 1938b. Noteworthy plants of southeastern Virginia. *Rhod.* 40:439-440.
- _____. 1950. *Gray's manual of botany*. 8th ed. New York: American Book Company.
- Gaiser, L.O. 1946. The genus *Liatris*. *Rhod.* 48:163-412.
- Goodman, George J. 1950. A new variety of *Saxifraga*. *Rhod.* 52:183.
- Gray, Fenton and H.M. Galloway. 1959. Soils of Oklahoma. *Okla. State Univ. Exp. Sta. Misc. Pub.* MP-56.
- Harlow, W.M. 1931. The identification of the pines of the United States, native and introduced, by needle structure. *New York State Coll. of Forestry Bul.* 4(2a):11-21.
- Hermann, Frederick J. 1946. The perennial species of *Utrica* in the United States east of the Rocky Mountains. *Am. Midl. Nat.* 35:773-778.
- Hitchcock, A.S. and Agnes Chase. 1950. *Manual of the grasses of the United States*. 2nd ed. U. S. Govt. Print. Off.
- Hitchcock, C. Leo. 1936. The genus *Lepidium* in the United States. *Madrona* 3:265-320.
- Hopkins, Milton. 1938. *Arabis* in eastern and central North America. *Rhod.* 39:63-76, 155-167, 175-179.
- _____. 1942. *Cercis* in North America. *Rhod.* 44:193-211.
- Hotchkiss, N. and H.L. Dozier. 1949. Taxonomy and distribution of the North American cattails. *Am. Midl. Nat.* 41:237-254.
- House, Homer D. 1908. The North American species of the genus *Ipomoea*. *Ann. N. Y. Acad. Sci.* 18:181-263.
- Isley, Duane. 1955. The Leguminosae of the North-Central United States II. Hedysareae. *Iowa State Coll. Jour. Sci.* 30(1):33-118.
- Kearney, T.H. 1955. A tentative key to the North American species of *Abutilon miller*. *Leaflet West. Bot.* 7(10):241-254.
- Larisey, Mary Maxine. 1940. A monograph on the genus *Baptisia*. *Ann. Mo. Bot. Gard.* 27:119-244.
- Mackenzie, K.K. 1931. Cyperaceae. *N. Am. Fl.* 18(1-7):1-478.
- Mathias, M.E. and L. Constance. 1945. Umbelliferae. *N. Am. Flora* 28b:43-295.
- Miser, Hugh D. 1954. Geologic map of the State of Oklahoma. *Dept. Inter. U. S. Geol. Surv.*
- Munz, Phillip A. 1965. Onagraceae. *N. Am. Flora II* 5:1-278.
- McClintock, Elizabeth and Carl Epling. 1942. A revision of the genus *Monarda* (Labiatae). *Univ. Calif. Publ. Bot.* 29(2):147-192.
- McCoy, Doyle. 1954. The genus *Lythrum* in Oklahoma. *Proc. Okla. Acad. Sci.* 33:156-158.
- McGregor, R.L. 1950. Two varieties of *Cystopteris fragilis*. *Am. Fern Jour.* 40:201-207.
- McKelvey, Susan Delano. 1955. Botanical exploration of the trans-Mississippi West 1790-1850. *Arn. Arb.*, Harvard Univ. 1144. pp.
- McVaugh, Roger. 1943. Campanulaceae (Lobelioideae). *N. Am. Flora* 32A:36-82.

- Ogden, E.C. 1945. The broad-leaved species of *Potamogeton* of North America north of Mexico. *Rhod.* 45:57-105, 119-163, 171-214.
- Ownbey, Gerald B. 1947. A monograph of the North American species of *Corydalis*. *An. Mo. Bot. Gard.* 34(3):187-252.
- Ownbey, Marion and Hannan C. Case. 1955. Cytotaxonomic studies in *Allium*. 1. The *Allium canadense* Alliance. *Research Studies of State Coll. of Wash. Monographic Sup.* 1:1-106.
- Palmer, E.J. 1924. The ligneous flora of Rich Mountain, Arkansas and Oklahoma. *Journ. Arn. Arb.* 5:108-134.
- _____. 1931. Conspectus of the genus *Amorpha*. *Journ. Arn. Arb.* 12:159-196.
- _____. 1932. Leaves from a collector's note book. *Journ. Arn. Arb.* 13:436.
- Payson, Edwin B. 1918. The North American species of *Aquilegia*. *Cont. U. S. Nat. Herbarium* 20(4):133-159.
- Pennell, Francis W. 1935. Scrophylariaceae of eastern temperate North America. *Acad. Nat. Sci. Phil. Monog.* 1.
- Perry, Lilly M. 1937. Variants in two species of *Dephinium* (*D. carolinianum* and *D. virescens*). *Rhod.* 39: 20-22.
- Perude, Robert E., Jr. 1957. Synopsis of *Rudbeckia* subgenus *Rudbeckia*. *Rhod.* 59:293-299.
- Russell, Norman H. 1965. Violets of central and eastern United States: An introductory survey. *Sida* 2(1):1-113.
- Rydberg, Per Axel. 1913. *Agrimonia*. *N. Am. Flora* 22(5):391-396.
- Sargent, C.S. 1922. *Manual of the trees of North America*. 2 vols. Dover Pub. Inc. N. Y.
- Shinners, Lloyd H. 1946. Revision of the genus *Kubnia*. *Wrightia* 1(2):122-144.
- _____. 1947. Revision of the genus *Krigia*. *Wrightia* 1(3):187-206.
- _____. 1951. *Agave lata*, a new species from North Texas and Oklahoma. *Field and Lab.* 19:171-173.
- Snider, L.C. 1917. Geography of Oklahoma. *Okla. Geol. Surv. Bul.* 27:23-325.
- Standley, Paul C. 1917. Amaranthaceae. *N. Am. Flora* 21(2):96-167.
- Stanford, E.E. 1926. *Polygonum hydropiperoides* and *Polygonum opelousanum*. *Rhod.* 28:23-27.
- Stevens, G.W. and C.W. Shannon. 1917. Plant life in Oklahoma. *Okla. Geol. Surv. Bul.* 27:215-246.
- Steyermark, Julian A. 1941. A study of *Arenaria patula*. *Rhod.* 43:325-333.
- Svenson, Henry Knut. 1957. *Eleocharis*. *N. Am. Flora* 18(9):509-540.
- Thwaites, Reuben G. 1905. Nuttall's travels into the Arkansa Territory, 1819. *Early western travels 1748-1846*. Vol 13:199-227.
- Turner, B.L. 1950. Vegetative key to Texas *Desmanthus* (Leguminosae) and similar genera. *Field and Lab.* 18(2):51-65.
- _____. 1951. Revision of the United States species of *Neptunia*. *Am. Midl. Nat.* 46:82-92.
- _____. 1955. The *Cassia fasciculata* complex (Leguminosae) in Texas. *Field and Lab.* 23(3-4):87-91.
- Tryon, R.M., Jr. 1941. A revision of the genus *Pteridium*. *Rhod.* 43:1-31, 37-67.
- Tryon, Alice F. 1957. A revision of the fern genus *Pellaea* section *Pellaea*. *Ann. Mo. Bot. Gard.* 44:125-148.
- Wahlgren, Harry F. 1941. Climate of Oklahoma. Climate and man. *U.S.D.A. Yearbook of Agri.* 1941:1065-1074.
- Waterfall, U.T. 1950. Some additions to the Oklahoma flora. *Rhod.* 52:35.
- _____. 1951. The genus *Callirhoe* (Malvaceae) in Texas. *Field and Lab.* 19(3):107-119.
- _____. 1954a. A new species of *Carex* (section *Phyllostachyae* from Oklahoma. *Rhod.* 56(661):21-23.
- _____. 1954b. Studies in the composition and distribution of the Oklahoma flora-XXI. *Rhod.* 56 (667):157-162.
- _____. 1966. *Keys to the flora of Oklahoma*. Stillwater Okla. Privately published.
- _____. 1967. *Physalis* in Mexico, Central America and the West Indies. *Rhod.* 69:82-329.
- Weatherby, C.A. 1927. The group of *Acalypha virginica* in eastern North America. *Rhod.* 29:193-204.

- Webber, John M. 1953. *Yuccas of the Southwest. U.S.D.A. Agri. Monograph* 17:1-73.
- Wheeler, L.C. 1941. *Euphorbia* subgenus *Chamaesyce* in Canada and the United States exclusive of southern Florida. *Rhod.* 43:97-154, 168-205, 223-286.
- Wherry, E.T. 1955. *The genus Phlox*. Philadelphia: Morris Arboretum.
- Wilbur, R.L. and H.S. Daoud. 1961. The genus *Lechea* (Cistaceae) in the southeastern United States. *Rhod.* 63:103-118.
- Woodson, Robert E., Jr. 1942. Commentary on the North American genera of Commelinaceae. *Ann. Mo. Bot. Gard.* 29:141-154.
- _____. 1954. The North American species of *Asclepias* L. *Ann. Mo. Bot. Gard.* 41(1):1-211.
- Yunkers, Truman G. 1943. Genus *Cuscuta*, Convolvulaceae. *Fl. of Texas* 3(2):123-150.

APPENDIX

Updated Flora of Southeastern Oklahoma
from the Sans Bois to the Kiamichi Mountains.

Editor's note: Originally this listing followed the Engler-Prantl system for families, as used in the *Keys to Flora of Oklahoma* (Waterfall 1966). Nomenclature has been revised according to the National Plant Data Center, Baton Rouge, LA (<http://plants.usda.gov>) and organized based on the Angiosperm Phylogeny Group, Missouri Botanical Gardens <http://www.mobot.org/MOBOT/research/APweb/>) Accessed December 2009. [EM]

FERNS

ASPLENIACEAE

Asplenium pinnatifidum Nutt.
Asplenium platyneuron (L.) Britton, Sterns & Poggenb.

DENNSTAEDTIACEAE

Pteridium aquilinum (L.) Kuhn var.
pseudocaudatum (Clute) A. Heller

DRYOPTERIDACEAE

Athyrium filix-femina (L.) Roth. ssp.
asplenioides (Michx.) Hultén
Cystopteris tennesseensis Shaver [syn =
Cystopteris fragilis var. *simulans*]
Dryopteris marginalis (L.) A. Gray
Polystichum acrostichoides (Michx.) Schott
Woodsia obtusa (Spreng.) Torr.

OSMUNDACEAE

Osmunda regalis L. var. *spectabilis* (Willd.)
Gray

POLYPODIACEAE

Pleopeltis polypodioides (L.) Andrews &
Windham ssp. *michauxiana* (Weath.)
Andrews & Windham

PTERIDACEAE

Adiantum pedatum L.
Cheilanthes lanosa (Michx.) D.C. Eaton [syn =
Cheilanthes vestita]
Pellaea atropurpurea (L.) Link

GYMNOSPERMS

CUPRESSACEAE

Juniperus virginiana L.
Taxodium distichum (L.) Rich.

PINACEAE

Pinus echinata Mill.
Pinus taeda L.

BASAL ANGIOSPERMS

ANNONACEAE

Asimina triloba (L.) Dunal

ARISTOLOCHIACEAE

Aristolochia tomentosa Sims

LAURACEAE

Lindera benzoin (L.) Blume var. *benzoin*
Sassafras albidum (Nutt.) Nees. [syn =
Sassafras albidum var. *molle*]

MAGNOLIACEAE

Magnolia acuminata (L.) L.

NYMPHACEAE

Brasenia schreberi J.F. Gmel.
Nelumbo lutea Willd.
Nuphar lutea (L.) Sm. ssp. *advena* (Aiton)
Kartesz & Gandhi [syn = *Nuphar advena*,
Nuphar advena var. *tomentosa*, *Nuphar*
ovata, *Nuphar ozarkana*]
Nymphaea odorata Aiton

SAURURACEAE

Saururus cernuus L.

MONOCOTS

AGAVACEAE

Manfreda virginica (L.) Salisb. ex Rose [syn =
Agave lata, *Agave virginica*]

ALISMATACEAE

Alisma plantago-aquatica L.
Echinodorus cordifolius (L.) Griseb.
Sagittaria ambigua J. G. Sm.
Sagittaria graminea Michx.
Sagittaria latifolia Willd.
Sagittaria platyphylla (Engelm.) J. G. Sm.

ARACEAE

Arisaema dracontium (L.) Schott.
Arisaema triphyllum (L.) Schott. ssp. *triphyllum*
[syn = *Arisaema atrorubens*]

COMMELINACEAE

Commelina communis L.
Commelina diffusa Burm. f.
Commelina erecta L. var. *angustifolia* (Michx.)
Fernald
Commelina erecta L. var. *deamiana* Fernald
Commelina erecta L. var. *erecta*
Commelina virginica L.
Tradescantia ernestiana E.S. Anderson &
Woodson
Tradescantia hirsuticaulis Small
Tradescantia hirsutiflora Bush
Tradescantia ohioensis Raf.
Tradescantia tharpia E.S. Anderson & Woodson

CYPERACEAE

Carex amphibola Steud.
Carex annectens (E.P. Bicknell) E.P. Bicknell
Carex bicknellii Britton
Carex blanda Dewey
Carex brevior (Dewey) Mack.
Carex bushii Mack. [syn = *Carex caroliniana*
var. *cuspidata*]
Carex cephalophora Muhl. ex Willd.
Carex crinita Lam. var. *brevicrinis* Fernald
Carex crus-corvi Shuttlw. ex Kunze
Carex flaccosperma Dewey

Carex frankii Kunth
Carex gravida L.H. Bailey var. *lunelliana*
(Mack) F.J. Herm.
Carex hyalina Boott
Carex jooii L.H. Bailey
Carex laevivaginata (Kük.) Mack.
Carex latebracteata Waterf.
Carex lupuliformis Sartwell ex Dewey
Carex lurida Wahlenb.
Carex meadii Dewey
Carex microrhyncha Mack.
Carex muhlenbergii Schkuhr ex Willd. var.
enervis Boott
Carex oklahomensis Mack. [syn = *Carex*
stipata var. *oklahomensis*]
Carex oxylepis Torr. & Hook.
Carex retroflexa Muhl. ex Willd.
Carex squarrosa L.
Carex stipata Muhl. ex Willd. var. *stipata*
Carex texensis (Torr.) L. H. Bailey
Carex tribuloides Wahlenb.
Carex vulpinoidea Michx.
Cyperus acuminatus Torr. & Hook. ex Torr.
Cyperus echinatus (L.) Alph. Wood [syn =
Cyperus ovularis var. *sphaericus*]
Cyperus erythrorhizos Muhl.
Cyperus lupulinus (Spreng.) Marcks ssp.
lupulinus
Cyperus strigosus L.
Cyperus virens Michx.
Eleocharis acicularis (L.) Roem. & Schult. var.
acicularis
Eleocharis compressa Sull. var. *acutisquamata*
(Buckley) S.G. Sm. [syn = *Eleocharis*
acutisquamata]
Eleocharis montevidensis Kunth
Eleocharis obtusa (Willd.) Schult.
Eleocharis quadrangulata (Michx.) Roem. &
Schult.
Fimbristylis autumnalis (L.) Roem. & Schult.
Fimbristylis dichotoma (L.) Vahl.
Fimbristylis thermalis S. Watson [syn =
Fimbristylis spadicea]
Fimbristylis vahlii (Lam.) Link.
Isolepis carinata Hook. & Arn. ex Torr. [syn =
Scirpus koilolepis]

Kyllinga brevifolia Rottb. [syn = *Cyperus brevifolius*]
Rhynchospora capitellata (Michx.) Vahl.
Rhynchospora corniculata (Lam.) A. Gray
Rhynchospora glomerata (L.) Vahl.
Rhynchospora harveyi Wm. Boott
Rhynchospora macrostachya Torr. ex A. Gray
Rhynchospora recognita (Gale) Kral [syn =
Rhynchospora globularis var. *recognita*]
Schoenoplectus americanus (Pers.) Volkart ex
Schinz & R. Keller [syn = *Scirpus americanus*]
Schoenoplectus californicus (C.A. Mey.) Palla
[syn = *Scirpus californicus*]
Scirpus atrovirens Willd.
Scirpus cyperinus (L.) Kunth
Scirpus lineatus Michx.

DIOSCOREACEAE

Dioscorea quaternata J.F. Gmel. [syn =
Dioscorea villosa var. *glabrifolia*]

IRIDACEAE

Iris cristata Aiton
Sisyrinchium angustifolium Mill.
Sisyrinchium campestre E.P. Bicknell

JUNCACEAE

Juncus acuminatus Michx.
Juncus brachycarpus Engelm.
Juncus bufonius L.
Juncus coriaceus Mack.
Juncus diffusissimus Buckley
Juncus effusus L. var. *solutus* Fernald &
Wiegand
Juncus interior Wiegand
Juncus marginatus Rostk.
Juncus repens Michx.
Juncus scirpoides Lam.
Juncus tenuis Willd.
Juncus validus Coville var. *validus* [syn =
Juncus crassifolius]
Luzula bulbosa (Alph. Wood) Smyth & Smyth
Luzula echinata (Small) F.J. Herm.

LILIACEAE

Aletris farinosa L.
Allium canadense L. var. *canadense*
Allium canadense L. var. *fraseri* Ownbey
Allium canadense L. var. *hyacinthoides* (Bush)
Ownbey & Aase
Allium canadense L. var. *mobile* (Regel)
Ownbey
Allium perdulce S.V. Fraser
Allium vineale L. ssp. *compactum* (Thuill.) Coss
& Germ.
Amianthium muscitoxicum (Walter) A. Gray
Camassia angusta (Engelm. & A. Gray) Blank.
Camassia scilloides (Raf.) Cory
Cooperia drummondii Herbert [syn =
Zephyranthes brazosensis]
Erythronium albidum Nutt.
Erythronium americanum Ker Gawl.
Hypoxis hirsuta (L.) Coville
Maiathemum racemosum (L.) Link ssp.
racemosum [syn = *Smilacina racemosa*
var. *cylindrata*]
Nothoscordum bivalve (L.) Britton
Polygonatum biflorum (Walter) Elliot var.
commutatum (Schult. & Schult. f.) Morong
[syn = *Polygonatum canaliculatum*]
Smilax bona-nox L.
Smilax glauca Walter
Smilax herbacea L.
Smilax tamnoides L.
Trillium viride Beck
Uvularia grandiflora Sm.
Veratrum woodii J.W. Robbins ex Alph. Wood
Yucca glauca Nutt.
Zigadenus nuttallii (A. Gray) S. Watson

LEMNACEAE

Lemna valdiviana Phil.
Spirodela polyrrhiza (L.) Schleid.

MARANTACEAE

Thalia dealbata Fraser ex Roscoe

NAJADACEAE

Najas guadalupensis (Spreng.) Magnus

ORCHIDACEAE

- Calopogon tuberosus* (L.) Britton, Sterns & Poggenb. var. *tuberosus* [syn = *Calopogon pulchellus*]
Spiranthes cernua (L.) Rich.
Spiranthes tuberosa Raf.
Spiranthes vernalis Engelm. & A. Gray

POACEAE

- Agrostis stolonifera* L. [syn = *Agrostis alba*]
Agrostis eliottiana Schult.
Agrostis hyemalis (Walt.) Britton, Sterns & Poggenb.
Agrostis perennans (Walt.) Tuck.
Aira elegans Willd. ex Kunth.
Alopecurus carolinianus Walter
Andropogon gerardii Vitman
Andropogon glomeratus (Walter) Britton, Sterns & Poggenb. var. *glomeratus* [syn = *Andropogon virginicus* var. *abbreviatus*]
Andropogon gyrans Ashe var. *gyrans* [syn = *A. eliottii*]
Andropogon ternarius Michx.
Andropogon virginicus L. var. *virginicus*
Aristida dichotoma Michx. var. *curtissii* Gray
Aristida dichotoma Michx. var. *dichotoma*
Aristida longespica Poir.
Aristida oligantha Michx.
Aristida purpurascens Poir.
Arundinaria gigantea (Walter) Muhl.
Axonopus festifolius (Raddi) Kuhl. [syn = *Axonopus affinis*]
Bothriochloa barbinodis Lag. [syn = *Andropogon barbinodis*]
Bothriochloa saccharoides (Sw.) Rydb. [syn = *Andropogon saccharoides*]
Bouteloua curtipendula (Michx.) Torr.
Bouteloua dactyloides (Nutt.) J.T. Columbus [syn = *Buchloe dactyloides*]
Brachyelytrum erectum (Schreb. ex Spreng.) P. Beauv.
Bromus arvensis L. [syn = *Bromus japonicus*]
Bromus catharticus Vahl
Bromus hordeaceus L. ssp. *hordeaceus* [syn = *Bromus mollis*]
Bromus inermis Leyss.
Bromus kalmii A. Gray [syn = *Bromus purgans*]

- Bromus secalinus* L.
Bromus tectorum L.
Cenchrus spinifex Cav. [syn = *Cenchrus incertus*, *Cenchrus pauciflorus*]
Chasmanthium latifolium (Michx.) Yates [syn = *Uniola latifolia*]
Chasmanthium laxum (L.) Yates [syn = *Uniola laxa*]
Chasmanthium sessiliflorum (Poir.) Yates [syn = *Uniola sessiliflora*]
Chloris verticillata Nutt.
Chloris virgata Sw.
Cinna arundinaceae L.
Coelorachis cylindrica (Michx.) Nash [syn = *Manisuris cylindrica*]
Cynosurus echinatus L.
Dactylis glomerata L.
Danthonia spicata (L.) P. Beauv. ex Roem. & Schult.
Diarrhena obovata (Gleason) Brandenburg [syn = *Diarrhena americana* var. *obovata*]
Dichantherium acuminatum (Sw.) Gould & C.A. Clark var. *fasciculatum* (Torr.) Freckmann [syn = *Panicum lanuginosum* var. *fasciculatum*]
Dichantherium acuminatum (Sw.) Gould & C.A. Clark var. *lindheimeri* (Nash) & C.A. Clark [syn = *Panicum lanuginosum* var. *lindheimeri*]
Dichantherium boscii (Poir.) Gould & C.A. Clark [syn = *Panicum boscii*]
Dichantherium dichotomum (L.) Gould var. *dichotomum* [syn = *Panicum dichotomum*]
Dichantherium laxiflorum (Lam.) Gould [syn = *Panicum laxiflorum*]
Dichantherium linearifolium (Scribn. ex Nash) Gould
Dichantherium malacophyllum (Nash) Gould [syn = *Panicum malacophyllum*]
Dichantherium oligosanthos (Schult.) Gould var. *scribnerianum* (Nash) Gould [syn = *Panicum oligosanthos* var. *helleri*, *Panicum oligosanthos* var. *scribnerianum*]
Dichantherium scoparium (Lam.) Gould [syn = *Panicum scoparium*]
Dichantherium sphaerocarpon (Elliot) Gould var. *isophyllum* (Scribn.) Gould & C.A.

- Clark [syn = *Panicum microcarpon*,
Panicum polyanthes]
Dichantherium sphaerocarpon (Elliot) Gould
 var. *sphaerocarpon* [syn = *Panicum*
sphaerocarpon]
Dichantherium villosissimum (Nash)
 Freckmann var. *praecocius* (Hitch. &
Chase) Freckmann [syn = *Panicum*
praecocius]
Dichantherium wilcoxianum (Vasey)
 Freckmann [syn = *Panicum wilcoxianum*]
Digitaria villosa (Walter) Pers. [syn = *Digitaria*
filiformis var. *villosa*]
Digitaria ischaemum (Schreb.) Schreb. ex
Muhl.
Digitaria sanguinalis (L.) Scop.
Digitaria violascens Link
Echinochloa colona (L.) Link
Echinochloa crus-galli (L.) P. Beauv.
Eleusine indica (L.) Gaertn.
Elymus canadensis L.
Elymus hystrix L. var. *hystrix* [syn = *Hystrix*
patula]
Elymus interruptus Buckley
Elymus submuticus (Hook.) Smyth & Smyth
 [syn = *Elymus virginicus* var. *submuticus*]
Elymus virginicus L. var. *virginicus* [syn =
Elymus virginicus var. *glabriflorus*, *Elymus*
virginicus var. *jejunus*]
Eragrostis capillaris (L.) Nees
Eragrostis frankii C.A. Mey. ex Steud.
Eragrostis hirsuta (Michx.) Nees
Eragrostis hypnoides (Lam.) Britton, Sterns &
Poggenb.
Eragrostis intermedia Hitchc.
Eragrostis japonica (Thunb.) Trin. [syn =
Eragrostis glomerata]
Eragrostis minor Host [syn = *Eragrostis*
poaeoides]
Eragrostis pilosa (L.) P. Beauv.
Eragrostis spectabilis (Pursh) Steud.
Eragrostis trichodes (Nutt.) Alph. Wood
Eriochloa contracta Hitchc.
Festuca paradoxa Desv.
Festuca subverticillata (Pers.) Alexeev [syn =
Festuca obtusa]
Gymnopogon ambiguus (Michx.) Britton,
Sterns & Poggenb.
Hordeum pusillum Nutt.
Leersia oryzoides (L.) Sw.
Leersia virginica Willd.
Leptochloa panicea (Retz) Ohwi ssp. *brachiata*
(Steud.) N. Snow [syn = *Leptochloa*
filiformis]
Lolium perenne L.
Lolium perenne L. ssp. *multiflorum* (Lam.)
Husnot [syn = *Lolium multiflorum*]
Melica mutica Walter
Muhlenbergia capillaris (Lam.) Trin.
Muhlenbergia sobolifera (Muhl. ex Willd.) Trin.
Muhlenbergia tenuiflora (Kunth.) Trin.
Panicum anceps Michx.
Panicum brachyanthum Steud.
Panicum capillare L.
Panicum dichotomiflorum Michx.
Panicum rigidulum Bosc ex Nees var. *rigidulum*
[syn = *Panicum agrostoides*]
Panicum virgatum L.
Paspalum dilatatum Poir.
Paspalum dissectum (L.) L.
Paspalum distichum L.
Paspalum floridanum Michx.
Paspalum laeve Michx.
Paspalum setaceum Michx.
Paspalum urvillei Steud.
Phalaris canariensis L.
Phalaris caroliniana Walter
Piptochaetium avenaceum (L.) Parodi [syn =
Stipa avenacea]
Poa annua L.
Poa pratensis L.
Poa sylvestris A. Gray
Saccharum brevibarbe (Michx.) Pers. var.
contortum (Elliot) R. Webster [syn =
Erianthus contortus]
Saccharum giganteum (Walter) Pers. [syn =
Erianthus giganteus]
Sacciolepis striata (L.) Nash
Schedonorus pratensis (Huds.) P. Beauv. [syn
= *Festuca elatior*]
Schizachyrium scoparium (Michx.) Nash var.
scoparium [syn = *Andropogon scoparius*]
Setaria italica (L.) P. Beauv.

Setaria parviflora (Poir.) Kerguélen [syn =
Setaria geniculata]
Setaria pumila (Poir.) Roem. & Schult. ssp.
pumila [syn = *Setaria glauca*]
Setaria viridis (L.) P. Beauv.
Sorghastrum nutans (L.) Nash
Sorghum halepense (L.) Pers.
Sphenopholis intermedia (Rydb.) Rydb.
Sphenopholis obtusata (Michx.) Scribn.
Sporobolus clandestinus (Biehler) Hitchc. [syn
= *Sporobolus asper* var. *canovirens*]
Sporobolus compositus (Poir.) Merr. var. *macer*
(Trin.) Kartesz & Gandhi [syn = *Sporobolus*
asper var. *macer*]
Sporobolus cryptandrus (Torr.) A. Gray
Sporobolus indicus (L.) R. Br. var. *indicus* [syn
= *Sporobolus poiretii*]
Sporobolus pyramidatus (Lam.) Hitchc.
Sporobolus vaginiflorus (Torr. ex A. Gray) Alph.
Wood var. *vaginiflorus*
Steinchisma hians (Elliot) Nash [syn = *Panicum*
hians]
Tridens flavus (L.) Hitchc.
Tridens strictus (Nutt.) Nash
Triplasis purpurea (Walter) Chapm.
Tripsacum dactyloides (L.) L.
Urochloa platyphylla (Munro ex C. Wright) R.D.
Webster [syn = *Brachiaria platyphylla*]
Vulpia myuros (L.) C.C. Gmel. [syn = *Festuca*
myuros]
Vulpia octoflora (Walter) Rydb. var. *octoflora*
[syn = *Festuca octoflora*]
Zizaniopsis miliacea (Michx.) Döll & Asch.

POTOMOGETONACEAE

Potamogeton diversifolius Raf.
Potamogeton pulcher Tuck.

SPARGANIACEAE

Sparganium americanum Nutt.

TYPHACEAE

Typha angustifolia L.
Typha domingensis Pers.
Typha latifolia L.

XYRIDACEAE

Xyris difformis Chapm.
Xyris torta Sm.

EUDICOTS

BERBERIDACEAE

Podophyllum peltatum L.

FUMARIACEAE

Corydalis crystallina Engelm.
Corydalis micrantha (Engelm. ex A. Gray) A.
Gray

MENISPERMACEAE

Calycocarpum lyonii (Pursh) A. Gray
Cocculus carolinus (L.) DC.

PAPAVERACEAE

Sanguinaria canadensis L.

PLATANACEAE

Platanus occidentalis L.

RANUNCULACEAE

Anemone berlandieri Pritz. [syn = *Anemone*
decapetala]
Anemone caroliniana Walter
Aquilegia canadensis L.
Clematis pitcheri Torr. & A. Gray
Clematis versicolor Small ex Rydb.
Delphinium carolinianum Walter
Delphinium tricornis Michx.
Delphinium wootonii Rydb. [syn = *Delphinium*
virescens]
Ranunculus abortivus L.
Ranunculus fascicularis Muhl. ex Bigelow
Ranunculus laxicaulis (Torr. & A. Gray) Darby
Ranunculus micranthus Nutt.
Ranunculus recurvatus Poir.
Thalictrum dasycarpum Fisch. & Avé-Lall.
Thalictrum thalictroides (L.) Spach. [syn =
Anemonella thalictroides]

ROSIDS**ACERACEAE**

Acer negundo L. var. *negundo*
Acer rubrum L.
Acer saccharum Marsh.

ANACARDIACEAE

Rhus aromatica Aiton var. *aromatica*
Rhus copallinum L. var. *latifolia* Engl.
Rhus glabra L.
Toxicodendron radicans (L.) Kuntze ssp.
radicans [syn = *Rhus radicans*]
Toxicodendron pubescens Mill. [syn = *Rhus toxicodendron*]

BETULACEAE

Alnus serrulata (Aiton) Willd.
Betula nigra L.
Carpinus caroliniana Walter
Ostrya virginiana (Mill.) K. Koch var. *virginiana*

BRASSICACEAE

Arabis canadensis L.
Arabis missouriensis Greene
Capsella bursa-pastoris (L.) Medik.
Cardamine concatenata (Michx.) Sw. [syn =
Dentaria laciniata]
Cardamine parviflora L. var. *arenicola* (Britton)
O. E. Schulz
Cardamine pensylvanica Muhl. ex Willd.
Draba brachycarpa Nutt. ex Torr. & A. Gray
Lepidium densiflorum Schrad.
Lepidium virginicum L.
Rorippa palustris (L.) Besser ssp. *fernaldiana*
(Butters & Abbe) Jonsell [syn = *Rorippa islandica* ssp. *fernaldiana*]
Selenia aurea Nutt.
Sibara virginica (L.) Rollins
Streptanthus maculatus Nutt.
Thlaspi arvense L.

CAPPARACEAE

Cleome spinosa Jacq.
Polanisia dodecandra (L.) DC. ssp.
trachysperma (Torr. & A. Gray) Iltis

CELASTRACEAE

Euonymus americanus L.
Euonymus atropurpureus Jacq.

CISTACEAE

Lechea mucronata Raf.
Lechea tenuifolia Michx.

CLUSIACEAE

Hypericum densiflorum Pursh
Hypericum lobocarpum Gattinger ex J.M.
Coult. [syn = *Hypericum densiflorum* var.
lobocarpum, *Hypericum oklahomense*]
Hypericum drummondii (Grev. & Hook.) Torr. &
A. Gray
Hypericum gentianoides (L.) Britton, Sterns &
Poggenb.
Hypericum hypericoides (L.) Crantz ssp.
hypericoides [syn = *Ascyrum hypericoides*]
Hypericum mutilum L.
Hypericum prolificum L. [syn = *Hypericum spathulatum*]
Hypericum pseudomaculatum Bush
Hypericum punctatum Lam.
Triadenum tubulosum (Walter) Gleason

CRASSULACEAE

Penthorum sedoides L.
Sedum nuttallianum Raf.

CUCURBITACEAE

Melothria pendula L.
Cucurbita foetidissima Kunth

EUPHORBIACEAE

Acalypha gracilens A. Gray [syn = *Acalypha gracilens* var. *fraseri*]
Acalypha monococca (Engelm. ex A. Gray) Lill.
W. Mill. & Gandhi [syn = *Acalypha gracilens* ssp. *monococca*]
Acalypha virginica L.
Chamaesyce maculata (L.) Small [syn =
Euphorbia supina]
Chamaesyce missurica (Raf.) Shinnery [syn =
Euphorbia missurica]
Chamaesyce nutans (Lag.) Small [syn =
Euphorbia nutans]

Chamaesyce prostrata (Aiton) Small [syn =
Euphorbia prostrata]
Cnidocolus texanus (Müll. Arg.) Small
Croton capitatus Michx. var. *capitatus*
Croton capitatus Michx. var. *lindheimeri*
(Engelm. and A. Gray) Müll. Arg.
Croton glanulosus L. var. *septentrionalis* Müll.
Arg.
Croton lindheimerianus Scheel
Croton michauxii G.L. Webster [syn =
Crotonopsis linearis]
Croton monanthogynus Michx.
Euphorbia cyathophora Murray [syn =
Euphorbia heterophylla var. *graminifolia*]
Euphorbia dentata Michx. var. *dentata*
Euphorbia pubentissima Michx. [syn =
Euphorbia corollata var. *paniculata*]
Euphorbia spathulata Lam. [syn = *Euphorbia*
obtusata]
Euphorbia tetrapora Engelm.
Leptopus phyllanthoides (Nutt.) G.L. Webster
[syn = *Andrachne phyllanthoides*]
Phyllanthus caroliniensis Walter
Stillingia sylvatica L.
Tragia betonicifolia Nutt.

FABACEAE

Amorpha fruticosa L. [syn = *Amorpha virgata*]
Amorpha laevigata Nutt.
Apios americana Medik.
Astragalus canadensis L.
Astragalus crassicaarpus Nutt. var. *crassicaarpus*
Astragalus crassicaarpus Nutt. var. *trichocalyx*
(Nutt.) Barneby
Astragalus distortus Torr. & A. Gray var.
distortus
Baptisia alba (L.) Vent. var. *macrophylla*
(Larisey) Isely [syn = *Baptisia leucantha*]
Baptisia australis (L.) R. Br. var. *minor* (Lehm.)
Fernald
Baptisia bracteata Muhl. ex Elliot var.
leucophaea (Nutt.) Kartesz & Gandhi [syn
= *Baptisia leucophaea*, *Baptisia*
leucophaea var. *glabrescens*]
Baptisia nuttalliana Small
Baptisia spaerocarpa Nutt.
Baptisia stricta Nutt.

Chamaecrista fasciculata (Michx.) Greene var.
fasciculata [syn = *Cassia fasciculata*,
Cassia fasciculata var. *rostrata*]
Chamaecrista nictitans (L.) Moench ssp.
nictitans var. *nictitans*
Cercis canadensis L. var. *canadensis*
Clitoria mariana L.
Dalea candida Michx. ex Willd. var. *candida*
Dalea purpurea Vent.
Desmanthus illinoensis (Michx.) MacMill. ex
B.L. Rob. & Fernald
Desmodium glutinosum (Muhl. ex Willd.) Alph.
Wood
Desmodium laevigatum (Nutt.) DC.
Desmodium marilandicum (L.) DC.
Desmodium nudiflorum (L.) DC.
Desmodium obtusum (Muhl. ex Willd.) DC. [syn
= *Desmodium rigidum*]
Desmodium perplexum B.G. Schub. [syn =
Desmodium paniculatum var. *dillenii*]
Desmodium paniculatum (L.) DC. var.
paniculatum
Desmodium sessilifolium (Torr.) Torr. & A.
Gray
Galactia volubilis (L.) Britt.
Gleditsia triacanthos L.
Kummerowia stipulacea (Maxim.) Makino [syn
= *Lespedeza stipulacea*]
Kummerowia striata (Thunb.) Schindl. [syn =
Lespedeza striata]
Lathyrus latifolius L.
Lathyrus pusillus Elliot
Lespedeza capitata Michx.
Lespedeza cuneata (Dum. Cours.) G. Don
Lespedeza hirta (L.) Hornem. ssp. *hirta*
Lespedeza procumbens Michx.
Lespedeza repens (L.) W. Bartram
Lespedeza stuevei Nutt. [syn = *Lespedeza*
stuevei var. *angustifolia*]
Lespedeza violacea (L.) Pers.
Lespedeza virginica (L.) Britt.
Mimosa microphylla Dryand. [syn = *Schrankia*
uncinata]
Neptunia lutea (Leavenworth) Benth.
Orbexilum pedunculatum (Mill.) Rydb. var.
pedunculatum [syn = *Psoralea psoralioides*
var. *eglandulosa*]

Orbexilum simplex (Nutt. ex Torr. & A. Gray)
Rydb. [syn = *Psoralea simplex*]
Psoralidium tenuiflora (Pursh) Rydb. [syn =
Psoralea tenuiflora]
Rynchosia latifolia Nutt. ex Torr. & A. Gray
Robinia pseudoacacia L.
Securigera varia (L.) Lassen [syn = *Coronilla*
varia]
Senna marilandica (L.) Link [syn = *Cassia*
marilandica]
Senna occidentalis (L.) Link [syn = *Cassia*
occidentalis]
Strophostyles helvola (L.) Elliot
Strophostyles leiosperma (Torr. & A. Gray)
Piper
Strophostyles umbellata (Muhl. ex Willd.)
Britton
Stylosanthes biflora (L.) Britton, Sterns &
Poggenb. [syn = *Stylosanthes biflora* var.
hispidissima]
Tephrosia onobrychoides Nutt.
Tephrosia virginiana (L.) Pers. [syn =
Tephrosia virginiana var. *holosericea*]
Trifolium arvense L.
Trifolium carolinianum Michx.
Trifolium dubium Sibth.
Trifolium incarnatum L.
Trifolium pratense L.
Trifolium reflexum L.
Vicia caroliniana Walter
Vicia minutiflora F.G. Dietr.

FAGACEAE

Castanea pumila (L.) Mill. var. *ozarkensis*
(Ashe) Tucker [syn = *Castanea ozarkensis*]
Fagus grandifolia Ehrh.
Quercus alba L.
Quercus coccinea Münchh.
Quercus falcata Michx. [syn = *Quercus falcata*
var. *triloba*]
Quercus lyrata Walter
Quercus macrocarpa Michx.
Quercus marilandica Münchh.
Quercus muehlenbergii Engelm.
Quercus nigra L.
Quercus pagoda Raf. [syn = *Quercus falcata*
var. *pagodifolia*]

Quercus palustris Münchh.
Quercus phellos L.
Quercus rubra L. var. *ambigua* (A. Gray)
Fernald [syn = *Quercus rubra* var. *borealis*]
Quercus shumardii Buckley var. *shumardii*
Quercus shumardii Buckley var. *schneckii*
(Britton) Sarg.
Quercus stellata Wangenh.
Quercus velutina Lam.

GERANIACEAE

Geranium carolinianum L.

GROSSULARIACEAE

Itea virginica L.
Ribes cynosbati L.

HALORAGACEAE

Myriophyllum aquaticum (Vell.) Verdc. [syn =
Myriophyllum brasiliense]
Myriophyllum heterophyllum Michx.
Myriophyllum pinnatum (Walt.) Britton, Sterns
& Poggenb.
Proserpinaca palustris L. var. *crebra* Fernald &
Grisc.

HAMAMELIDACEAE

Hamamelis vernalis Sarg.
Liquidambar styraciflua L.

HIPPOCASTANACEAE

Aesculus glabra Willd.

JUGLANDACEAE

Carya alba (L.) Nutt. [syn = *Carya tomentosa*]
Carya aquatica (Michx. f.) Nutt.
Carya cordiformis (Wangenh.) K. Koch
Carya illinoensis (Wangenh.) K. Koch
Carya myristiciformis (Michx. f.) Nutt.
Carya ovata (Mill.) K. Koch
Carya texana Buckley
Juglans nigra L.

LINACEAE

Linum medium (Planch.) Britton var. *texanum*
(Planch.) Fernald

Linum sulcatum Riddell

LYTHRACEAE

Didiplis diandra (Nutt. ex DC.) Alph. Wood [syn = *Peplis diandra*]

Lythrum alatum Pursh var. *alatum*

Rotala ramosior (L.) Koehne

MALVACEAE

Abutilon theophrasti Medik.

Callirhoe alcaeoides (Michx.) A. Gray

Callirhoe pedata (Nutt. ex Hook.) A. Gray [syn = *Callirhoe digitata* var. *stipulata*]

Hibiscus lasiocarpus Cav.

Hibiscus laevis All. [syn = *Hibiscus militaris*]

Malva pusilla L. [syn = *Malva rotundifolia*]

Sida rhombifolia L.

Sida spinosa L.

MELASTOMATACEAE

Rhexia mariana L. var. *interior* (Pennell) Kral & Bostick [syn = *Rhexia interior*]

MELIACEAE

Melia azedarach L.

MORACEAE

Maclura pomifera (Raf.) C.K. Schneid.

Morus alba L.

Morus rubra L.

ONAGRACEAE

Gaura longiflora Spach [syn = *Gaura filiformis*]

Gaura sinuata Nutt. ex Ser.

Ludwigia alternifolia L.

Ludwigia decurrens Walter [syn = *Jussiaea decurrens*]

Ludwigia glandulosa Walter ssp. *glandulosa*

Ludwigia palustris (L.) Elliot

Ludwigia peploides (Kunth) P.H. Raven ssp. *peploides* [syn = *Jussiaea peploides*]

Oenothera elata Kunth. ssp. *hirsutissima* (A. Gray ex S. Watson) W. Dietr. [syn = *Oenothera biennis* var. *hirsutissima*]

Oenothera fruticosa L.

Oenothera laciniata Hill

Oenothera linifolia Nutt.

Oenothera speciosa Nutt.

Oenothera villosa Thunb. ssp. *villosa* [syn = *Oenothera biennis* var. *canescens*]

OXALIDACEAE

Oxalis corniculata L.

Oxalis stricta L.

Oxalis violacea L.

PASSIFLORACEAE

Passiflora incarnata L.

Passiflora lutea L.

POLYGALACEAE

Polygala incarnata L.

Polygala polygama Walter

Polygala sanguinea L.

Polygala verticillata L., var. *isocycla* Fernald

RHAMNACEAE

Berchemia scandens (Hill.) K. Koch

Ceanothus americanus L.

Ceanothus herbaceus Raf. var. *pubescens* (T. & G.) Shinnars

Frangula caroliniana (Walter) A. Gray [syn = *Rhamnus caroliniana*]

ROSACEAE

Agrimonia parviflora Aiton

Agrimonia pubescens Wallr.

Agrimonia rostellata Wallr.

Amelanchier arborea (Michx. f.) Fernald

Crataegus crus-galli L.

Crataegus marshallii Ettl.

Crataegus pruinosa (Wendl. f.) Koch. [syn = *Crataegus mackenziei*]

Crataegus punctata Jacq. [syn = *Crataegus collina*]

Crataegus spathulata Michx.

Crataegus uniflora Münchh.

Crataegus viridis L.

Geum canadense Jacq. var. *canadense*

Geum canadense Jacq. var. *texanum* Fernald & Weath.

Gillenia stipulata (Muhl. ex Willd.) Baill.

Potentilla simplex Michx.

Prunus americana Marsh., var. *americana*
Prunus angustifolia Marsh.
Prunus mexicana S. Watson
Prunus munsoniana W. Wright & Hedrick
Prunus serotina Ehrh.
Rosa carolina L.
Rosa foliolosa Nutt. ex Torr. & A. Gray
Rosa setigera Michx. var. *setigera*
Rosa setigera Michx. var. *tomentosa* Torr. & A. Gray
Rubus aboriginum Rydb.
Rubus argutus Link. [syn = *Rubus louisianus*]
Rubus bushii L.H. Bailey [syn = *Rubus ozarkensis*, *Rubus scibilis*]
Rubus oklahomus L.H. Bailey
Rubus trivialis Michx.
Sanguisorba annua (Nutt. ex Hook.) Nutt. ex Torr. & A. Gray

RUTACEAE

Ptelea trifoliata L. ssp. *trifoliata*

SALICACEAE

Populus deltoides Bartram ex Marsh.
Salix caroliniana Michx.
Salix humilis Marsh. var. *humilis*
Salix interior Rowlee
Salix nigra Marsh.

SAPINDACEAE

Sapindus saponaria L. var. *drummondii* (Hook. & Arn.) L.D. Benson [syn = *Sapindus drummondii*]

SAXIFRAGACEAE

Heuchera americana L. var. *americana*
Saxifraga texana Buckley

STAPHYLEACEAE

Staphylea trifolia L.

TILIACEAE

Tilia americana L.
Tilia americana L. var. *americana* [syn = *Tilia neglecta*]

ULMACEAE

Celtis laevigata Willd.

Celtis occidentalis L.
Celtis tenuifolia Nutt.
Ulmus alata Michx.

URTICACEAE

Boehmeria cylindrica (L.) Sw.
Laportea canadensis (L.) Weddell
Parietaria pennsylvanica Muhl. ex Willd.
Pilea pumila (L.) A. Gray

VISCACEAE

Phoradendron leucarpum (Raf.) Reveal & M.C. Johnst. [syn = *Phoradendron serotinum*]

VITACEAE

Ampelopsis arborea (L.) Koehne
Ampelopsis cordata Michx.
Cissus trifoliata (L.) L.
Parthenocissus quinquefolia (L.) Planch
Vitis acerifolia Raf.
Vitis aestivalis Michx.
Vitis cinerea (Engelm.) Engelm. ex Millard
Vitis rotundifolia Michx.
Vitis rupestris Scheele
Vitis vulpina L.

ZYGOPHYLLACEAE

Tribulus terrestris L.

ASTERIDS**ACANTHACEAE**

Dicliptera brachiata (Pursh) Spreng.
Justicia americana (L.) Vahl
Ruellia humilis Nutt.
Ruellia pedunculata Torr. ex A. Gray
Ruellia strepens L.

AMARANTHACEAE

Amaranthus albus L. [syn = *Amaranthus graecizans*]
Amaranthus retroflexus L.
Amaranthus spinosus L.
Froelichia gracilis (Hook.) Moq.
Iresine rhizomatosa Standl.

APIACEAE

- Ammoselinum butleri* (Engelm. ex S. Watson)
J.M. Coult. & Rose
Chaerophyllum tainturieri Hook. var. *tainturieri*
[syn = *Chaerophyllum texanum*]
Cicuta maculata L.
Cryptotaenia canadensis (L.) DC.
Cynosciadium digitatum DC.
Daucus pusillus Michx.
Eryngium prostratum Nutt. ex DC.
Eryngium yuccifolium Michx. var. *synchaetum*
A. Gray ex J.M. Coult. & Rose
Hydrocotyle verticillata Thunb.
Limnosciadium pinnatum (DC.) Mathias &
Constance
Osmorhiza longistylis (Torr.) DC.
Polytaenia nuttallii DC.
Ptilimnium capillaceum (Michx.) Raf.
Ptilimnium nuttallii (DC.) Britton
Sanicula canadensis L.
Spermolepis echinata (Nutt. ex DC.) A. Heller
Spermolepis inermis (Nutt. ex DC.) Mathias &
Constance
Thaspium barbinode (Michx.) Nutt.
Trepocarpus aethusae Nutt. ex DC.
Zizia aurea (L.) W.D.J. Koch

APOCYNACEAE

- Amsonia illustris* Woodson
Amsonia tabernaemontana Walter
Apocynum cannabinum L.
Trachelospermum difforme (Walter) A. Gray

AQUIFOLIACEAE

- Ilex decidua* Walter
Ilex opaca Aiton

ASCLEPIADACEAE

- Asclepias amplexicaulis* Sm.
Asclepias hirtella (Pennell) Woodson
Asclepias obovata Elliot
Asclepias quadrifolia Jacq.
Asclepias syriaca L.
Asclepias tuberosa L.
Asclepias verticillata L.
Asclepias viridiflora Raf. [syn = *Asclepias*
viridiflora var. *lanceolata*]

- Asclepias viridis* Walter
Matelea baldwyniana (Sweet) Woodson
Matelea gonocarpos (Walter) Shinnars

ASTERACEAE

- Achillea millefolium* L. var. *occidentalis* DC.
[syn = *Achillea lanulosa*]
Ageratina altissima (L.) King & H. Rob. var.
altissima [syn = *Eupatorium rugosum*]
Ambrosia artemisiifolia L. var. *elatior* (L.)
Descourtils
Ambrosia bidentata Michx.
Ambrosia psilostachya DC. [syn = *Ambrosia*
psilostachya var. *lindheimeriana*]
Ambrosia trifida L. var. *texana* Scheele
Antennaria plantaginifolia (L.) Richardson
Anthemis cotula L.
Arctium minus Bernh.
Arnoglossum plantagineum Raf. [syn = *Cacalia*
plantaginea]
Astranthium integrifolium (Michx.) Nutt.
Baccharis halimifolia L.
Bidens aristosa (Michx.) Britton [syn = *Bidens*
polylepis, *Bidens aristosa* var. *mutica*]
Bidens bipinnata L.
Bidens discoidea (Torr. & A. Gray) Britton
Bidens frondosa L.
Boltonia asteroides (L.) L'Hér. var. *latisquama*
(A. Gray) Cronquist
Boltonia asteroides (L.) L'Hér. var. *recognita*
(Fernald & Grisc.) Cronquist
Boltonia diffusa Elliot
Brickellia eupatorioides (L.) Shinnars var.
texana (Shinnars) Shinnars [syn = *Kuhnia*
eupatorioides var. *ozarkana*]
Centaurea americana Nutt.
Chaetopappa asteroides Nutt. ex DC.
Chrysopsis pilosa Nutt.
Cirsium altissimum (L.) Hill
Cirsium carolinianum (Walter) Fernald & B.G.
Schub.
Conoclinium coelestinum (L.) DC. [syn =
Eupatorium coelestinum]
Conyza canadensis (L.) Cronquist var.
canadensis
Conyza canadensis (L.) Cronquist var. *glabrata*
(A. Gray) Cronquist

- Coreopsis grandiflora* Hogg ex Sweet var. *grandiflora*
Coreopsis grandiflora Hogg ex Sweet var. *harveyana* (A. Gray) Sherff
Coreopsis palmata Nutt.
Coreopsis tinctoria Nutt. var. *tinctoria*
Coreopsis tripteris L.
Crepis pulchra L.
Croptilon divaricatum (Nutt.) Raf. [syn = *Haplopappus divaricatus*]
Echinacea angustifolia DC. var. *angustifolia*
Echinacea angustifolia DC. var. *strigosa* R.L. McGregor
Echinacea pallida (Nutt.) Nutt.
Echinacea purpurea (L.) Moench
Eclipta prostrata (L.) L. [syn = *Eclipta alba*]
Elephantopus carolinianus Raeusch.
Erechtites hieracifolia (L.) Raf. ex DC.
Erigeron pulchellus Michx.
Erigeron strigosus Muhl. ex. Willd.
Erigeron tenuis Torr. & A. Gray
Eupatorium perfoliatum L.
Eupatorium serotinum Michx.
Eurybia hemispherica (Alexander) G.L. Nesom [syn = *Aster hemisphericus*]
Euthamia gymnospermoides Greene [syn = *Solidago gymnospermoides*]
Facelis retusa (Lam.) Sch. Bip. [syn = *Facelis apiculata*]
Fleischmannia incarnata (Walter) King & H. Rob. [syn = *Eupatorium incarnatum*]
Gaillardia aestivalis (Walter) H. Rock var. *aestivalis* [syn = *Gaillardia lanceolata* var. *fastigiata*, *Gaillardia serotina*]
Gamochaeta purpurea (L.) Cabrera [syn = *Gnaphalium purpureum*]
Grindelia lanceolata Nutt.
Helenium amarum (Raf.) H. Rock var. *amarum*
Helenium flexuosum Raf.
Helianthus angustifolius L.
Helianthus annuus L.
Helianthus hirsutus Raf. [syn = *Helianthus hirsutus* var. *trachyphyllus*, *Helianthus hirsutus* var. *stenophyllus*]
Helianthus mollis Lam.
Heliopsis helianthoides (L.) Sweet var. *scabra* (Dunal) Fernald
Heterotheca subaxillaris (Lam.) Britton & Rusby [syn = *Heterotheca latifolia*]
Hieracium gronovii L.
Hieracium longipilum Torr.
Hymenopappus scabiosaeus L'Hér. var. *scabiosaeus*
Krigia caespitosa (Raf.) K.L. Chambers [syn = *Krigia oppositifolia*]
Krigia dandelion (L.) Nutt.
Krigia occidentalis Nutt.
Krigia virginica (L.) Willd.
Lactuca canadensis L. [syn = *Lactuca canadensis* var. *latifolia*]
Lactuca serriola L. [syn = *Lactuca scariola*]
Liatris aspera Michx. var. *aspera*
Liatris aspera Michx. var. *intermedia* (Lunell) Gaiser
Liatris elegans (Walter) Michx.
Liatris mucronata DC.
Liatris pycnostachya Michx.
Liatris squarrosa (L.) Michx. var. *glabrata* (Rydb.) Gaiser
Liatris squarrosa (L.) Michx. var. *hirsuta* (Rydb.) Gaiser
Liatris squarrosa Michx. [syn = *Liatris scabra*]
Marshallia caespitosa Nutt. ex DC.
Mikania scandens (L.) Willd.
Oligoneuron nitidum (Torr. & A. Gray) Small [syn = *Solidago nitida*]
Pakera obovata (Muhl. ex Willd.) W.A. Weber & A. Löve [syn = *Senecio obovatus* var. *rotundus*]
Pakera tomentosa (Michx.) C. Jeffrey [syn = *Senecio tomentosus*]
Parthenium integrifolium L.
Pluchea camphorata (L.) DC.
Pityopsis graminifolia (Michx.) Nutt. var. *tenuifolia* (Torr.) Semple & F.D. Bowers [syn = *Chrysopsis microcephala*]
Polymnia canadensis L.
Prenanthes altissima L.
Pseudognaphalium obtusifolium (L.) Hilliard & B.L. Burt ssp. *obtusifolium* [syn = *Gnaphalium obtusifolium*]
Pyrrhopappus grandiflorus (Nutt.) Nutt. [syn = *Pyrrhopappus scaposus*]

Rudbeckia grandiflora (D. Don) J.F. Gmel. ex DC.
Rudbeckia hirta L. var. *pulcherrima* Farw.
Rudbeckia maxima Nutt.
Rudbeckia subtomentosa Pursh
Rudbeckia triloba L.
Silphium asteriscus L.
Silphium laciniatum Torr. var. *robinsonii* L.M. Perry
Smallanthus uvedalius (L.) Mack. ex Small [syn = *Polymnia uvedalia* var. *densipilis*]
Solidago altissima L. [syn = *Solidago canadensis* var. *scabra*]
Solidago caesia L.
Solidago canadensis L. var. *gilvocanescens* Rydb.
Solidago hispida Muhl. ex Willd.
Solidago missouriensis Nutt. var. *fasciculata* Holz.
Solidago nemoralis Aiton
Solidago odora Aiton
Solidago petiolaris Aiton
Solidago radula Nutt.
Solidago rugosa Mill. ssp. *aspera* (Aiton) Cronquist
Solidago speciosa Nutt. var. *rigidiscula* Torr. & A. Gray [syn = *Solidago speciosa* var. *angustata*]
Solidago ulmifolia Muhl. ex Willd. var. *microphylla* A. Gray [syn = *Solidago delicatula*]
Sonchus asper (L.) Hill
Symphotrichum anomalum (Engelm.) G.L. Nesom [syn = *Aster anomalus*]
Symphotrichum cordifolium (L.) G.L. Nesom [syn = *Aster sagittifolius*]
Symphotrichum ericoides (L.) G.L. Nesom var. *ericoides* [syn = *Aster ericoides*]
Symphotrichum lateriflorum (L.) A. Löve & D. Löve var. *lateriflorum* [syn = *Aster lateriflorus*]
Symphotrichum oolentangiense (Riddell) G.L. Nesom var. *oolentangiens* [syn = *Aster azureus*]
Symphotrichum patens (Aiton) G.L. Nesom var. *patentissimum* (Lindl. ex DC.) G.L.

Nesom [syn = *Aster patens* var. *patentissimum*]
Symphotrichum praealtum (Poir.) G.L. Nesom var. *praealtum* [syn = *Aster praealtus*]
Symphotrichum turbinellum (Lindl.) G.L. Nesom [syn = *Aster turbinellus*]
Taraxacum laevigatum (Willd.) DC. [syn = *Taraxacum erythrospermum*]
Verbesina alternifolia (L.) Britton ex Kearney [syn = *Actinomeris alternifolia*]
Verbesina encelioides (Cav.) Benth. & Hook. f. ex A. Gray
Verbesina helianthoides Michx.
Verbesina virginica L.
Vernonia baldwinii Torr. ssp. *baldwinii*
Vernonia fasciculata Michx.
Vernonia gigantea (Walter) Trel. ssp. *gigantea* [syn = *Vernonia altissima*]
Vernonia lettermannii Engelm. ex A. Gray
Vernonia missurica Raf.
Xanthium strumarium L.

BALSAMINACEAE

Impatiens capensis Meerb.

BIGNONIACEAE

Campsis radicans (L.) Seem. ex Bureau
Catalpa bignonioides Walter

BORAGINACEAE

Cynoglossum virginianum L.
Hackelia virginiana (L.) I.M. Johnst.
Heliotropium indicum L.
Lithospermum incisum Lehm.
Myosotis verna Nutt.

BUDDLEJACEAE

Polypremum procumbens L.

CACTACEAE

Opuntia ficus-indica (L.) Mill. [syn = *Opuntia compressa*]
Opuntia macrorhiza Engelm. var. *macrorhiza* [syn = *Opuntia tortispina*]

CALLITRICHACEAE

Callitriche heterophylla Pursh

CAMPANULACEAE

- Lobelia appendiculata* A. DC.
Lobelia cardinalis L.
Lobelia puberula Michx.
Lobelia spicata Lam. var. *leptostachys* (A. DC.)
 Mack. and Bush
Triodanis biflora (Ruiz & Pav.) Greene [syn =
Specularia biflora]
Triodanis lamprosperma McVaugh [syn =
Specularia lamprosperma]
Triodanis leptocarpa (Nutt.) Nieuwl. [syn =
Specularia leptocarpa]
Triodanis perfoliata (L.) Nieuwl. [syn =
Specularia perfoliata]

CAPRIFOLIACEAE

- Lonicera flava* Sims
Lonicera japonica Thunb.
Lonicera sempervirens L.
Sambucus nigra L. ssp. *canadensis* (L.) R. Bolli
Symphoricarpos orbiculatus Moench
Viburnum rufidulum Raf. [syn = *Viburnum*
prunifolium var. *ferrugineum*]

CARYOPHYLLACEAE

- Cerastium brachypodium* (Engelm. ex A. Gray)
 B.L. Rob.
Cerastium fontanum Buamg. ssp. *vulgare*
 (Hartm.) Greuter & Burdet [syn =
Cerastium vulgatum]
Cerastium glomeratum Till. [syn = *Cerastium*
viscosum]
Minuartia drummondii (Shinners) McNeill [syn =
Arenaria drummondii]
Minuartia patula (Michx.) Mattf. [syn = *Arenaria*
patula]
Paronychia fastigiata (Raf.) Fern.
Paronychia virginica Spreng.
Sagina decumbens (Elliot) Torr. & A. Gray
Silene antirrhina L.
Silene stellata (L.) W.T. Aiton
Silene virginica L.
Stellaria media (L.) Vill.

CHENOPODIACEAE

- Chenopodium album* L.

- Chenopodium ambrosioides* L. var.
ambrosioides
Chenopodium pumilio R. Br.
Monolepis nuttalliana (Schult.) Greene

CONVOLVULACEAE

- Convolvulus arvensis* L.
Ipomoea hederacea Jacq.
Ipomoea lacunosa L.
Ipomoea pandurata (L.) G. Mey.
Ipomoea purpurea (L.) Roth.
Ipomoea quamoclit L.

CORNACEAE

- Cornus drummondii* C.A. Mey.
Cornus florida L.
Cornus obliqua Raf.
Nyssa sylvatica Marsh.

CUSCUTACEAE

- Cuscuta compacta* Juss. ex Choisy
Cuscuta cuspidata Engelm.
Cuscuta glomerata Choisy
Cuscuta indecora Choisy
Cuscuta pentagona Engelm. var. *glabrior*
 (Engelm.) Gandhi, R.D. Thomas & S.L.
 Hatch [syn = *Cuscuta glabrior*]
Cuscuta pentagona Engelm. var. *pentagona*
 [syn = *Cuscuta campestris*]

DROSERACEAE

- Drosera brevifolia* Pursh. [syn = *Drosera*
annua]

EBENACEAE

- Diospyros virginiana* L. [syn = *Diospyros*
virginiana var. *pubescens*]

ERICACEAE

- Lyonia ligustrina* (L.) DC. var. *foliosiflora*
 (Michx.) Fernald [syn = *Lyonia ligustrina*
 var. *salicifolia*]
Rhododendron oblongifolium (Small) Millais
Vaccinium arboreum Marsh.
Vaccinium pallidum Aiton [syn = *Vaccinium*
vacillans]
Vaccinium stamineum L.
Vaccinium virgatum Aiton

GENTIANACEAE

Gentiana saponaria L.
Sabatia angularis (L.) Pursh
Sabatia campestris Nutt.

HYDRANGEACEAE

Hydrangea arborescens L.

HYDROPHYLLACEAE

Hydrolea ovata Nutt. ex Choisy
Hydrolea uniflora Raf.
Nemophila phacelioides Nutt.
Phacelia glabra Nutt.
Phacelia hirsuta Nutt.

LAMIACEAE

Agastache nepetoides (L.) Kuntze
Blephilia ciliata (L.) Benth.
Cunila origanoides (L.) Britton
Hedeoma hispida Pursh
Lamium amplexicaule L.
Lycopus americanus Muhl. ex W. Bartram [syn =
= *Lycopus americanus* var. *scabrifolius*]
Lycopus rubellus Moench
Lycopus virginicus L.
Marrubium vulgare L.
Mentha spicata L.
Monarda fistulosa L. ssp. *fistulosa* var. *fistulosa*
Monarda punctata L. ssp. *punctata* var.
villicaulis (Pennell) Palmer & Steyerl.
Monarda russeliana Nutt. ex Sims [syn =
Monarda virgata]
Perilla frutescens (L.) Britt.
Physostegia angustifolia Fernald
Physostegia intermedia (Nutt.) Engelm. & A.
Gray
Physostegia virginiana (L.) Benth.
Prunella vulgaris L. ssp. *lanceolata* (W.
Bartram) Hultén
Pycnanthemum albescens Torr. & A. Gray
Pycnanthemum tenuifolium Schrad.
Salvia azurea Michx. ex Lam. var. *grandiflora*
Benth.
Salvia lyrata L.
Scutellaria elliptica Muhl. ex Spreng.
Scutellaria ovata Hill
Stachys palustris Nutt. var. *pilosa*

Stachys tenuifolia Willd.

Teucrium canadense L. var. *canadense* [syn =
Teucrium canadense var. *virginicum*]

Trichostema brachiatum L.

LENTIBULARIACEAE

Utricularia gibba L. [syn = *Utricularia biflora*]

LOGANIACEAE

Mitreola petiolata (J.F. Gmel.) Torr. & A. Gray
[syn = *Cynoctonum mitreola*]

Spigelia marilandica (L.) L.

MOLLUGINACEAE

Mollugo verticillata L.

NYCTAGINACEAE

Mirabilis albida (Walter) Heimerl

Mirabilis nyctaginea (Michx.) MacMill.

OLEACEAE

Chionanthus virginicus L.

Fraxinus americana L.

Fraxinus pennsylvanica Marsh.

PHYTOLACCACEAE

Phytolacca americana L.

PLANTAGINACEAE

Plantago aristata Michx.

Plantago lanceolata L.

Plantago rhodosperma Decne.

Plantago rugelii Decne.

Plantago virginica L.

POLEMONIACEAE

Ipomopsis rubra (L.) Wherry [syn = *Gilia rubra*]

Phlox cuspidata Scheele

Phlox pilosa L.

POLYGONACEAE

Brunnichia ovata (Walter) Shinnery [syn =
Brunnichia cirrhosa]

Eriogonum longifolium Nutt. var. *longifolium*
[syn = *Eriogonum vespinum*]

Polygonum convolvulus L.

Polygonum hydropiper L.

Polygonum hydropiperoides Michx. [syn =
Polygonum hydropiperoides var.
bushmanum, *Polygonum hydropiperoides*
var. *opelousanum*]

Polygonum orientale L.

Polygonum pensylvanicum L.

Polygonum persicaria L.

Polygonum punctatum Elliot

Polygonum ramosissimum Michx.

Polygonum scandens L. var. *scandens*

Polygonum tenue Michx.

Polygonum virginianum L. [syn = *Tovara*
virginiana]

Rumex acetosella L.

Rumex altissimus Alph. Wood

Rumex crispus L.

Rumex hastatulus Baldw.

Rumex pulcher L.

PORTULACACEAE

Claytonia virginica L.

Portulaca halimoides L. [syn = *Portulaca*
parvula]

Portulaca oleracea L.

Phemeranthus parviflorum (Nutt.) Kiger [syn =
Talinum parviflorum]

PRIMULACEAE

Dodecatheon meadia L.

Hottonia inflata Elliot

Lysimachia lanceolata Walter

RUBIACEAE

Cephalanthus occidentalis L. var. *occidentalis*

Diodia teres Walter var. *teres* [syn = *Diodia*
teres var. *setifera*]

Diodia virginiana L.

Galium aparine L.

Galium arkansanum A. Gray

Galium pilosum Aiton var. *pilosum*

Houstonia longifolia Gaertn. [syn = *Hedyotis*
purpurea var. *longifolia*]

Houstonia purpurea L. var. *purpurea* [syn =
Hedyotis purpurea]

Houstonia pusilla Schoepf [syn = *Hedyotis*
crassifolia]

Mitchella repens L.

Stenaria nigricans (Lam.) Terrell var. *nigricans*
[syn = *Hedyotis nigricans*]

SAPOTACEAE

Sideroxylon lanuginosum Michx. ssp.

lanuginosum [syn = *Bumelia lanuginosa*]

SCROPHULARIACEAE

Agalinis fasciculata (Elliot) Raf. [syn = *Gerardia*
fasciculata]

Agalinis gattingeri (Small) Small [syn =
Gerardia gattingeri]

Agalinis tenuifolia (Vahl.) Raf. var. *parviflora*
(Nutt.) Pennell [syn = *Gerardia tenuifolia*
ssp. *parviflora*]

Aureolaria grandiflora (Benth.) Pennell var.
grandiflora [syn = *Gerardia grandiflora*]

Aureolaria pectinata (Nutt.) Pennell [syn =
Gerardia pectinata]

Bacopa rotundifolia (Michx.) Wettst.

Buchnera americana L.

Castilleja coccinea (L.) Spreng.

Castilleja indivisa Engelm.

Collinsia violacea Nutt.

Gratiola brevifolia Raf.

Gratiola virginiana L.

Lindernia dubia (L.) Pennell var. *anagallidea*
(Michx.) Cooperr. [syn = *Lindernia*
anagallidea]

Lindernia dubia (L.) Pennell var. *dubia*

Mecardonia acuminata (Walter) Small var.
acuminata [syn = *Bacopa acuminata*]

Mimulus alatus Aiton

Nutallanthus texanus (Scheele) D.A. Sutton
[syn = *Linaria canadensis* var. *texana*]

Pedicularis canadensis L. ssp. *canadensis* [syn
= *Pedicularis canadensis* var. *dobbsii*]

Penstemon arkansanus Pennell

Penstemon digitalis Nutt. ex Sims

Penstemon tubiflorus Nutt.

Scrophularia marilandica L.

Verbascum blattaria L.

Verbascum thapsus L.

Veronica arvensis L.

Veronica peregrina L. ssp. *peregrina*

Veronicastrum virginicum (L.) Farw.

SOLANACEAE

Datura stramonium L.
Physalis angulata L. [syn = *Physalis angulata*
var. *lanceifolia*, *Physalis angulata* var.
pendula]
Physalis cinerascens (Dunal) Hitch. var.
cinerascens [syn = *Physalis viscosa* var.
cinerascens]
Physalis cordata Mill.
Physalis heterophylla Nees
Physalis pubescens L. var. *integrifolia* (Dunal)
Waterf.
Physalis pumila Nutt.
Physalis turbinata Medik.
Physalis virginiana Mill. var. *virginiana*
Solanum carolinense L.
Solanum elaeagnifolium Cav.
Solanum nigrum L.
Solanum rostratum Dunal

STYRACACEAE

Halesia carolina L.

VALERIANACEAE

Valerianella longiflora (Torr. & A. Gray) Walp.
Valerianella nuttallii (Torr. & A. Gray) Walp.
Valerianella radiata (L.) Dufur. [syn =
Valerianella stenocarpa var. *parviflora*]

VERBENACEAE

Callicarpa americana L.

Glandularia canadensis (L.) Nutt. [syn =
Verbena canadensis]
Glandularia pumila (Rydb.) Umber [syn =
Verbena pumila]
Phryma leptostachya L.
Verbena bracteata Cav. ex Lag. & Rodr.
Verbena halei Small
Verbena stricta Vent.
Verbena urticifolia L.

VIOLACEAE

Viola bicolor Pursh [syn = *Viola kitaibeliana*
var. *rafinisquei*]
Viola langloisii Greene, nom. inq.
Viola × *lovelliana* Brainerd (pro sp.)
[*missouriensis* × *triloba*]
Viola missouriensis Greene
Viola pedata L. [syn = *Viola pedata* var.
lineariloba]
Viola pubescens Aiton var. *pubescens* [syn =
Viola pensylvanica]
Viola pubescens Aiton var. *scabruiscula*
Swein. ex Torr. & A. Gray [syn = *Viola*
pensylvanica var. *leiocarpon*]
Viola × *primulifolia* L. (pro sp.) [*lanceolata* ×
macloskeyi]
Viola sagittata Aiton
Viola sororia Willd. [syn = *Viola papilionacea*]
Viola triloba Schwein. var. *dilatata* (Elliot)
Brainerd
Viola villosa Walter